

Assessment of Quality of Life of HIV Positive Patients Attending Anti-Retroviral Clinic in Enugu State University Teaching Hospital, Enugu, Nigeria

Ogbodo OC^{1*}, and Ossai EN²

¹Department of Community Medicine and Primary Care, Enugu State University College of Medicine, Parklane, Enugu, Nigeria

²Department of Community Medicine, University of Nigeria, Enugu Campus

***Corresponding author:** Ogbodo OC, Department of Community Medicine and Primary Care, Enugu State University College of Medicine, Parklane, Enugu, Nigeria.

Submitted: 14 May 2024 **Accepted:** 22 May 2024 **Published:** 31 May 2024

Citation: Ogbodo, O. C., & Ossai, E. N. (2024). Assessment of Quality of Life of HIV Positive Patients Attending Anti-Retroviral Clinic in Enugu State University Teaching Hospital, Enugu, Nigeria. *Journal of Sexual Health and AIDS Research*, 1(3), 01-12.

 Crossref Doi: 10.63620/MKJSHAR.2024.1006

Abstract

Introduction: Globally, more than 70 million people have been infected with the HIV virus and about 35 million people have died of HIV related causes since 1981.

Objective: To assess the quality of life of HIV positive patients attending Anti-retroviral clinic in Enugu State University Teaching Hospital, Enugu, Nigeria.

Methodology: This cross-sectional study was carried out among 400 respondents selected by systematic sampling. **Results:** Majority (75%) of the respondents were female, and more than half (54.3%) were more than 38 years old. 77.5% perceived their overall quality of life to be in excellent. Those 38 years and above had the best quality of life, while those within the age group 18 to 22 years had the worst quality of life. Married or widowed respondents had better excellent quality of health than those who were separated or single. The quality of life of majority of the respondents was affected by lack of money (75.8%), their physical condition (24.3%), psychological feelings (19.8%), social relationship (11.5%), and stigmatization (8.5%). Also, most of the respondents (56.2%) who were ill had the least quality of life, while those who were not ill (87.5%) had better quality of life. Majority of the respondents (68.3%) were satisfied with their physical condition. While 75.8% were satisfied with their psychological feelings. Also, majority of the respondents were satisfied with their ability to perform daily activities (78.8%), with social relationship with other people (77.5%), and with the environment of where they were living (78.5%).

Conclusion and Recommendations: The overall quality of life of majority of the respondents was good, and majority of them were satisfied with their current health status. There should be improvement on early diagnosis and treatment, including social support services.

Keywords: HIV/AIDS, Quality of life, Anti-retroviral Treatment

Introduction

The first Human immunodeficiency virus infection (HIV) case was reported in USA in 1981 [1]. Since then HIV infection has spread globally [2]. The population groups at increased risk of HIV transmission include: men who have sex with men, people who inject drugs, people in prisons and other closed settings, sex workers and their clients, and transgender people [2]. HIV infection drastically affects the patients' physical condition, socio-cultural relations, mental health, and economic aspects of life [3]. However, Anti-retroviral treatment (ART) has changed

the course of HIV infection from a rapidly progressive catastrophic illness to a chronic disease with reduction in mortality rate and prolongation of life [4].

More than 70 million people have been infected with the HIV virus and about 35 million people have died of HIV related causes since 1981 [2]. Sub-Saharan Africa bears most of the global HIV/AIDS burden [2]. Nigeria ranks second in sub-Saharan Africa in HIV/AIDS burden with a prevalence of is 2.9% among the adult population (aged 15-49 years) in 2016[5].

The Quality of Life (QoL) of HIV positive patients usually improve when they are receiving treatment at an ART clinic, and could further encourage the patient to continue their treatment [6]. Therefore, assessment of Quality of life has become an outcome measure in the management of HIV positive patients in ART clinics. This will help to evaluate the human and financial cost and benefits of new programs and interventions [7,8].

Methodology

The study site was the antiretroviral (ARV) clinic at Enugu State University Teaching Hospital, Enugu, Enugu State Nigeria. This antiretroviral clinic is accessed mostly by people living in Enugu State, Nigeria. Enugu is the capital of Enugu State located in the Southeast geo-political zone of Nigeria. The people living in Enugu State are predominantly Igbo speaking tribe and mainly Christians. They are mainly farmers, traders and civil servants. The estimated population of Enugu state in 2017 based on the 2006 Nigeria's census is 4,126,227 [9].

Study Design

This study was a facility-based cross-sectional analytical study.

Study Participants

The study participants were HIV positive patients who are age 18 years and older attending anti-retroviral clinic in Enugu State University Teaching Hospital, Enugu, Enugu State, Nigeria.

Sample Size Determination

The minimum sample size was determined by the formula for cross sectional study used for single proportions [10]. A total of 400 participants were used for the study based on the proportion of 79.75% that had excellent overall quality of life in a previous study, and on a type 1 error (α) of 0.05 [11].

Sampling Technique

The eligible and consenting adult patients were selected by systematic sampling technique. The last three months attendance of adult patients prior to this study was calculated using the facility's register. In the last three months prior to this study, the number of active adult male patients per month was 517, while the number of active adult female patients per month was 1,498. Therefore, the ratio of male adults to female adult patients was approximately 1:3. The study sample size of 400 was propor-

tionately allocated to male and female patients in the ratio of 1:3. Thus 100 male patients and 300 female patients were used for this study. The number of active male patients (517), and active female patients (1,498) were used as the sampling frame for the male and female patients respectively. The sampling interval was determined by dividing the sampling frames with the sample size 100, and 300 for male and female patients respectively.

Study Instruments

A pretested, semi-structured, interviewer-administered questionnaire adapted from WHOQol-HIV brief instrument and WHOQoL-HIV instrument users' manual was used to collect data on socio-demographic factors, quality of life, and satisfaction with the services at the antiretroviral clinic [12, 13].

Outcome Measure

The percentage Quality of Life (QoL) scores for each domain in the WHOQol-HIV brief instrument was calculated as the sum of individual scores obtained in the domain divided by the total attainable score in that domain multiply by 100. The factors that affects quality of life of the participants were determined by using tables, frequencies, proportions, and cross tabulations to determine the effect of socio-economic, and health related factors on the QoL of the respondents. Satisfaction level was assessed by the proportion of the participants who were satisfied with the services they received at the antiretroviral clinic.

Data Analysis

Data were collected and edited manually same day to detect omissions and to maintain uniform coding. The SPSS statistical package, version 21 was used for data entry and analysis. For descriptive variables, data was presented using table, frequencies and percentages, means, and cross tabulations. Categorical variables were analysed by use of proportions, while continuous variables that included QoL scores were summarised through the use of means and standard deviation. Testing between proportions was carried out using Fisher Exact and chi square test where appropriate. Student t-test was used for the analysis of statistical differences between the mean scores of QoL for dichotomous variables. Spearman's correlation was calculated to assess the effect of certain factors on mean QoL scores of the respondents. The level of statistical significance was set at $p < 0.05$.

Results

Table 1: Socio-demographic characteristics of the respondents

Variable	Frequency (n=400)	Percent (%)
Gender		
Male	100	25.0
Female	300	75.0
Age (years)		
Overall Mean (\pm SD)	39.3 \pm 12.3	
Mean age (male) (\pm SD)	43.8 \pm 13.9	
Mean age (female) (\pm SD)	37.8 \pm 11.4	
Age of respondents in groups		
18-22	17	4.3
23-27	47	11.8

28-32	76	19.0
33-37	43	10.8
38 and above	217	54.3
Tribe		
Ibo	351	87.8
Yoruba	13	3.3
Hausa	25	6.3
Other	11	2.8
Religion		
Christianity	374	93.5
Islam	25	6.3
African Traditional Religion	1	0.3
Education		
None	21	5.2
Primary	115	28.8
Secondary	179	44.8
Post-secondary	85	21.3
Marital status		
Single	93	23.3
Married	279	69.8
Living as married	3	0.8
Separated	2	0.5
Divorced	23	5.8
Employment status ((respondents)		
Unemployed	72	18.0
Self employed	250	62.5
Salary employed	78	19.5
Employment status (spouse/partner)		
No spouse/partner	147	36.8
Unemployed	23	5.5
Self employed	184	45.3
Salary employed	50	12.5

Table 1 shows the socio-demographic characteristics of the respondents. The overall mean age of the respondents was 39.3 years. The mean age of the males was 43.8 years, while that of the females was 37.8 years. There is a high literacy level among the respondents as only 5.2% of them did not attempt any formal

education. Majority of the respondents were married (69.8%), and only a few (5.8%) were divorced. Majority of the respondents (62.5%) were self-employed. However, 43.3% of the respondents' spouse/partner were self-employed, while 36.8% of the respondents did not have a spouse/partner.

Table 2: Clients health status

Variable	Frequency (n=400)	Percent (%)
General health condition		
Very poor	6	1.5
Poor	14	3.5
Neither Poor nor Good	34	8.5
Good	306	76.5
Very Good	40	10.0
Currently ill		
No	272	68.0
Yes	128	32.0

WHO Clinical Stage		
Stage 1	294	73.5
Stage 2	81	20.3
Stage 3	20	5.0
Stage 4	5	1.3
Perceived mode of HIV infection		
Sex with a man	96	24.0
Sex with a woman	31	7.8
Injection of drugs	24	6.0
Blood products	51	12.8
Sharp objects/ health facility related	31	7.8
Could not tell how	165	41.3
Mother to child	2	0.5

Table 2 shows the health status of the respondents at the time of the study. Majority of the respondents (76.5%) perceived their health status to be good, 68% of them were not ill at the time of their participation in this study. Most of the respondents (73.5%) were in WHO HIV clinical stage 1. A good proportion (41.3%)

of the respondents did not know how they became infected with HIV virus. The total proportion of the respondents that perceived their mode of infection to be through sexual intercourse was 31.8% (24% of males and 7.8% of females).

Table 3: Overall Quality of Life (QoL) of the respondents.

Variables	Frequency (n=400)	Percent (%)
Overall quality of life		
Poor	29	7.3
Good	61	15.3
Excellent	310	77.5

Table 3 shows the overall quality of life of the respondents. Majority (77.5%) of the respondents perceived their health to be in excellent condition, while 15.3% of the respondents perceived

their health to be in good condition. A low percentage (7.3%) of the respondents were in poor health.

Table 4: General factors influencing respondents' overall quality of life

Variable	Frequency n=400	Percent (%)
Physical condition	97	24.3
Psychological feelings	79	19.8
Level of independence	19	4.8
Social relationship	46	11.5
Environment	31	7.8
Lack of information	21	5.3
Lack of money	303	75.8
Your spirituality	21	5.3
Stigmatization	34	8.5

Table 4 shows the general factors influencing respondents' overall quality of life

The quality of life of majority of the respondents was affected by lack of money (75.8%), their physical condition (24.3%), psychological feelings (19.8%), social relationship (11.5%), and stigmatization (8.5%).

Table 5: Demographic and economic factors influencing respondents' overall quality of life.

Demographic Factor	Overall quality of life			Total	Chi-square (P-value)
	Poor n=29	Good n=61	Excellent n=310		
Gender					
Males	5(5.0)	13(13.0)	82(82.0)	100(100)	1.72(0.42)
Females	24(8.0)	48(16.0)	228(76.0)	300(100)	
Marital status					
Single	10(10.5)	22(23.7)	63(65.6)	93(100)	0.01FT
Married	18(6.4)	33(11.8)	230(81.6)	279(100)	
Separated	0(0)	0(0)	3(75.0)	3(100)	
Divorced	0(0)	2(100)	0(0)	2(100)	
Widowed	1(4.3)	4(17.4)	18(78.3)	23(100)	
Age					
18-22	3(17.6)	5(29.4)	9(52.9)	17(100)	0.02FT
23-27	3(6.4)	11(23.4)	33(70.2)	47(100)	
28-32	8(10.5)	12(15.8)	56(73.7)	76(100)	
33-37	1(2.3)	10(23.3)	32(74.4)	43(100)	
38 and above	14(6.5)	23(10.6)	180(82.9)	217(100)	
Educational					
None	2(9.5)	4(19.0)	15(71.4)	21(100)	0.58FT
Primary	6(5.2)	14(12.2)	95(82.6)	115(110)	
Secondary	12(6.7)	31(17.3)	136(76.0)	179(100)	
Tertiary	9(10.6)	12(14.1)	64(75.3)	85(100)	

Table 5 shows the demographic and economic factors influencing respondents' overall quality of life. Gender and educational level of the respondents did not influence the quality of life of the respondents. Married or widowed respondents had better ex-

cellent quality of health than those who were separated or single. The respondents that are 38 years or above had the best quality of life, while those within the age group 18 to 22 years had the worst quality of life.

Table 6: Health factors influencing the overall quality of life

Demographic Factor	Overall quality of life			Total	Chi-square (P-value)
	Poor n=29	Good n=61	Excellent n=310		
Perceived health					
Very poor	5(83.3)	0(0)	1(16.7)	6(100)	0.00FT
Poor	6(42.9)	4(28.6)	4(26.7)	14(100)	
Neither	6(17.1)	17(50.0)	11(32.4)	34(100)	
Good	12(3.9)	36(11.8)	258(84.3)	306(100)	
Very good	0(0)	4(10.0)	36(90.0)	40(100)	
Consider yourself ill					
No	9(3.3)	25(9.2)	238(87.5)	272(100)	49.64(0.00)
Yes	20(15.6)	36(28.1)	72(56.2)	128(100)	
Clinical staging					
Stage 1	22(7.5)	46(15.6)	226(76.9)	297(100)	0.83FT
Stage 2	5(6.2)	14(17.3)	62(76.5)	81(100)	
Stage 3	2(10)	1(5.0)	17(85.0)	20(100)	
Stage 4	0(0)	0(0)	5(100)	5(100)	

Table 6 shows the health factors influencing the overall quality of life. Most of the respondents (56.2%) who were ill had the

least quality of life, while those who were not ill (87.5%) had better quality of life.

Table 7: Respondents' level of satisfaction with their health

Variable	Frequency (n=400)	Percent (%)
Physical condition	8	2.0
Very dissatisfied	16	4.0
Dissatisfied	21	5.3
Neither satisfied nor dissatisfied	273	68.3
Satisfied	82	20.5
Very satisfied		
Psychological feeling	13	3.3
Very dissatisfied	14	3.5
Dissatisfied	39	9.8
Neither satisfied nor dissatisfied	303	75.8
Satisfied	31	7.8
Very satisfied		
Level of independence		
Very dissatisfied	13	3.3
Dissatisfied	9	2.3
Neither satisfied nor dissatisfied	44	11.0
Satisfied	315	78.8
Very satisfied	19	4.8
Social relationship		
Very dissatisfied	14	3.5
Dissatisfied	6	1.5
Neither satisfied nor dissatisfied	45	11.3
Satisfied	310	77.5
Very Satisfied	25	6.3
Satisfaction with their Environment		
Very dissatisfied	15	3.8
Dissatisfied	14	3.5
Neither satisfied nor dissatisfied	48	12.0
Satisfied	314	78.5
Very satisfied	9	2.3

Table 7 shows respondents' level of satisfaction with their health. Majority of the respondents (68.3%) were satisfied with their physical condition. While 75.8% were satisfied with their psychological feelings. Also, majority of the respondents were satisfied with their ability to perform daily activities (78.8%), with social relationship with other people (77.5%), and with the environment of where they were living (78.5%).

Discussion

Majority of the respondents were females (75%) [14, 15, 16]. This is similar to other findings in studies done in Lagos, Nigeria (61%), Uyo, South South, Nigeria (60%), in Kogi state, Nigeria, and in Ho Municipality, Ghana (73.4%) [17]. The highest proportion (54.3%) of the total respondents in this study was within

age group 38 and above years. This suggests that age group (38 and above) is an important risk group in HIV epidemic in Enugu State, Nigeria.

The mean age of the respondents was 39.3 ± 12.3 years. However, the average age of the males was 43.8 years, while that of the females was 37.8 years. This is similar to the mean age of respondents (38.1 ± 9 years) in a similar study done in Ibadan, Nigeria: Quality of life of People living with HIV and AIDS attending the Antiretroviral Clinic, University College Hospital, Nigeria [18]. Majority of the respondents were married (69.8%), while 23.3% of the respondents were single. Similarly, most of the respondents in a similar study done in an ART clinic in Lagos, Nigeria were married (61.4%) [14].

Majority (77.5%) of the respondents' quality of life was excellent. This could be as a result of the fact that majority of the respondents (73.5%) were in WHO HIV clinical stage 1, and high proportion of them (68%) were not ill at the time of this study. Over half of the participants (56%) in a multi-ethnic study rated their QoL as 'good or very good' [19]. In contrast, in United Kingdom, people living with HIV have significantly lower QoL than do the general population [20]. QoL of HIV positive patients is also significantly low in Xin-jiang, West China. In Cross River State, Nigeria, majority of the respondents rated their QoL scores as good (46.3%), very good (17.1%) [21]. Also, majority of the patients (84.7%) attending a HIV clinic in Lagos, Nigeria perceived their quality of life to be good [14]. However, significant proportion of HIV positive patients on ART in Uyo, Nigeria (78.5%) perceived their overall QoL to be poor [15].

The respondents' perception of their QoL regarding different aspects of health was varied. Majority (61.8%) of the respondents' physical condition was poor. This could be attributed to loss of weight associated with HIV infection. Also, mentally and physically, HIV positive patients attending ART clinics are not as healthy as their healthy neighbours [17]. In UK, physical health mean scores for HIV patients receiving treatment was higher than scores in other health domains [22]. But in India, the quality of life score was lowest for physical condition of the HIV patients receiving treatment [23]. In other aspects of health, a high proportion of the respondents (75.8%) in this study had good psychological feeling, 72% of the respondents consider their relationship with their environment to be good, 60% had excellent level of independence, almost half (49%) of the patients had excellent social relationship, while 92.5% of them had very poor level of spirituality.

Similar respondents in a study done in India reported highest QoL score in spirituality/religion/personal belief domain than in physical, psychological, social, and level of independence domains [24]. In contrast, in Ibadan, Nigeria, Kwara State Nigeria, the social domain recorded the lowest mean score [18, 25]. In Sao Paulo Brazil, worst average scores were reported in the Environment and Level of independence domains, while in Burkina Faso, the Environment and Level of independence domains also had lower scores compared to other quality of life domains [26, 27]. In Kogi state, north-central Nigeria, respondents that participated in similar study reported lower scores in the environmental and social domains [16].

Gender and educational factors did not influence the quality of life of the respondents in this study. However, similar studies done in Nigeria showed that gender influenced the quality of life of the respondents [16, 21, 18]. In Kogi State, Nigeria, female HIV patients have a higher QoL score when compared to their male counterparts in all domains [16]. Studies done in Cross River State, and Ibadan, in Nigeria did not report any significant difference in QoL scores between males and females living with HIV/AIDS and receiving care at ART clinics [21, 18].

Married or widowed respondents in this study had better excellent quality of health than those who were separated or single. Similarly, in Ilorin, Kwara State, Nigeria, HIV sero-positive married women have the highest QoL scores in all the domains compared to those with a different marital status. Also, in Uyo,

South South Nigeria, HIV positive patients who are single, separated or widowed have poor QoL in social and environmental domains [25, 28]. Respondents who are 38 years or above had the best quality of life, while those within the age group 18 to 22 years had the worst quality of life. Similarly, in China, and Iran, older age is also associated with a high QoL score [29, 30]. In contrast, in Portugal, older people (40 years and above) living with HIV have lower QoL in a number of domains (physical, level of independence, and social relationships) [31]. QoL of HIV patients in Croatia is also improved by being of younger age [32]. Also, in United States of America, younger age is associated with better QoL among HIV positive patients [33].

In increasing magnitude, stigmatization, social relationship, psychological feelings, physical condition of the patient, and lack of money reduces the QoL of the respondents in this study. Respondents in this study with better health had high quality of life, while those who were ill had the least quality of life. This may be because HIV patients with poor medical history are inclined to have worse QoL [29]. This can also explain why respondents in similar study done in India, 66 reported better daily routine activities (level of independence) and social activities in asymptomatic patients compared to those with AIDS defining symptoms [34]. Similarly, in Bangladesh, it was observed that asymptomatic HIV patients have better QoL that permits them to still perform their normal activity [35]. Among HIV positive patients on ART in Uganda, number of visits to clinic, level of education, WHO HIV stage and level of depression are determinants of physical health score [36]. Also, symptomatic HIV patients in Ghana, and Nigeria, significantly presented with a lower overall quality of life [17, 18].

In Brazil, having acquired opportunistic infections were predictors associated with a poorer quality of life [37]. It has also been observed in various studies across the world, that stigmatization lowers QoL scores [38-40]. Social support is also positively associated with the QoL of HIV/AIDS patients [41-43]. It has been observed in India, and Brazil, that presence and severity of symptoms are associated with lower physical domain score and overall QoL of HIV patients [24, 26]. In Ghana, HIV patients' self-appraisal of their health significantly predicted their quality of life, with lower QoL recorded among those who perceived themselves as ill [17]. In United States of America, younger age, higher income, and better social relation and support are associated with better QoL among HIV positive patients [33].

A high proportion (68.3%) of the respondents in this study was satisfied with their physical condition. Also, 75.8% were satisfied with their psychological feelings, 78.8% were satisfied with their ability to perform daily activities, and 77.5% were satisfied with social relationship with other people. A high proportion (78.5%) of the respondents was also satisfied with the environment of where they were living. This is because most of the respondents in this study are in WHO HIV stage 1, and were not suffering from serious HIV co-infection diseases. However, remarkably, in Vietnam, the proportions of respondents completely satisfied with overall service quality and treatment outcomes at HIV clinics were 42.4% and 18.8%, respectively [44]. Also, in Southern Ethiopia, 46.4% of HIV positive patients attending ART clinic were satisfied with the services they received [45]. Similar study done in Cross River, Nigeria, showed that majority

of the respondents (48%) were 'satisfied' with their health [21, 46-113].

Conclusion

Majority of the adult patients attending antiretroviral clinic at Enugu State University Teaching Hospital were females (75%) of the respondents were female, while over half (54.3%) of them were aged between ages 38 and above. Majority of the respondents were married (69.8%), and 23.3% of the respondents were single. The overall quality of life of majority of the respondents was good, and majority of them were satisfied with their current health status. This could be attributed to the fact that majority of the respondents were in WHO HIV clinical stage 1, and because only few of them were sick at the time of the study.

Majority of the respondents (68.3%) were satisfied with their physical condition. While 75.8% were satisfied with their psychological feelings. Also, majority of the respondents were satisfied with their ability to perform daily activities (78.8%), with social relationship with other people (77.5%), and with the environment of where they were living (78.5%). However, a very high proportion of the respondents (92.5%) were bothered by people blaming them, fear for their future, and worry about death. Age, marital status, and the current health of the respondents were the factors that determined the QoL of the respondents.

Recommendations

Government/Policy Makers and Health Workers

1. Government, policy makers, and health care workers should focus more on early diagnosis and treatment of HIV patients. This will ensure a better QoL of for these patients when they are receiving treatment.
2. They should also encourage enrolment of males at the ART clinics by health education, and counselling at every contact with them.

Health Workers

Health workers should improve or adjust their social support services available to these patients. This will positively improve their spirituality, religious, and personal beliefs.

Ethical Approval

Ethical approval and informed consent process for the study was obtained from Enugu State University Teaching Hospital ethical committee. Permission for this study was obtained from Enugu state ministry of health through the Enugu state ministry of health ethical committee on research projects. Permission was also obtained from the management of Enugu State University Teaching Hospital. Informed consent was obtained from each participant and participation in the study was voluntary. All information from this study was kept confidential and the information was used in such a way that no individual who participated in the study was linked to any information. The respondents were given the opportunity to withdraw from the study at any time during the study without any consequences to them.

Limitations

This study is a cross sectional study and therefore cannot draw conclusions about causality.

References

1. Centers for Disease Control and Prevention. (2001). First reports of AIDS. Morbidity and Mortality Weekly Reports, 50(21), 429-431, 439.
2. World Health Organization. (2017). Global Health Observatory data: HIV/AIDS [Internet]. Geneva. Retrieved from <http://www.who.int/gho/hiv/en/>
3. Abrefa-Gyan, T., Cornelius, L. J., & Okundaye, J. (2016). Socio-demographic factors, social support, quality of life, and HIV/AIDS in Ghana. *Journal of Evidence-Informed Social Work*, 13(2), 206-216.
4. Préau, M., Lepout, C., Salmon-Céron, D., Carrieri, P., Portier, H., & دغل اطم مورگ APROCO-COPILOTE. (2004). Health-related quality of life and patient-provider relationships in HIV-infected patients during the first three years after starting PI-containing antiretroviral treatment. *AIDS Care*, 16(6), 649-661.
5. The Joint United Nations Programme on HIV/AIDS (UNAIDS). (2017). Country fact sheet- Nigeria 2016 [Internet]. Geneva. Retrieved from <http://www.unaids.org/en/region-scountries/countries/nigeria>
6. Jin, Y., Liu, Z., Wang, X., Liu, H., Ding, G., Zhang, Y., & Lu, H. (2014). A systematic review of cohort studies of the quality of life in HIV/AIDS patients after antiretroviral therapy. *International Journal of STD & AIDS*, 25(10), 771-777.
7. Levine, S., & Croog, S. (1998). What constitutes quality of life? A conceptualization of the dimensions of life quality in healthy population and patients with cardiovascular disease. In N. Wenger, M. E. Mattson, C. D. Furberg, & J. Elinson (Eds.), *Assessment of quality of life in clinical trials of cardiovascular therapies* (pp. 46-58). Le Jacq.
8. Robinson, J. R., Young, T. K., Roos, L. L., & Gelskey, D. E. (1997). Estimating the burden of disease. Comparing administrative data and self-reports. *Medical Care*, 35(9), 932-947.
9. National Population Commission of Nigeria. (2006). Nigeria National Population Census 2006. Abuja.
10. Katz, D. L., Elmore, J. G., Wild, D. M., & Lucan, S. C. (2014). *Jekel's epidemiology, biostatistics, preventive medicine, and public health* (4th ed.). Elsevier Saunders.
11. Cummins, R. A. (1997). *Comprehensive Quality of Life Scale-Adult. Manual* (5th ed.). School of Psychology, Deakin University.
12. World Health Organization. (2002). WHOQOL-HIV BREF. Geneva.
13. World Health Organization. (2002). WHOQOL-HIV instrument. Geneva.
14. Olagunju, A. T., Ogundipe, O. A., Olagunju, T. O., & Adeyemi, J. D. (2013). A multi-dimensional assessment of quality of life among attendees of a West African HIV clinic and its use in tracking outcome. *HIV & AIDS Review*, 12(2), 63-67.
15. Motilewa, O. O., Ekanem, U. S., Onayade, A., & Sule, S. S. (2015). A Comparative Study of Health Related-Quality of Life Among HIV Patients on Pre-HAART and HAART in Uyo South-South Nigeria. *Journal of Antivirals & Antiretrovirals*, 7(3), 60-68.
16. Fatiregun, A. A., Mofolorunsho, K. C., & Osagbemi, K. G. (2009). Quality of life of people living with HIV/AIDS in Kogi State, Nigeria. *Benin Journal of Postgraduate Medicine*, 11(1), 21-27.

17. Osei-Yeboah, J., Owiredun, W. K. B. A., Norgbe, G. K., Lokpo, S. Y., Obirikorang, C., & et al. (2017). Quality of Life of People Living with HIV/AIDS in the Ho Municipality, Ghana: A Cross-Sectional Study. *AIDS Research and Treatment*, 2017, 6806951.
18. Folasire, O. F., Irabor, A. E., & Folasire, A. M. (2012). Quality of life of people living with HIV and AIDS attending the antiretroviral clinic, university college hospital, Nigeria. *African Journal of Primary Health Care & Family Medicine*, 4(1), 1-8.
19. Catalan, J., Tuffrey, V., Ridge, D., & Rosenfeld, D. (2017). What influences quality of life in older people living with HIV? *AIDS Research and Therapy*, 14(1), 22.
20. Miners, A., Phillips, A., Kreif, N., Rodger, A., Speakman, A., Collins, S., & Rabkin, M. (2014). Health-related quality-of-life of people with HIV in the era of combination antiretroviral treatment: A cross-sectional comparison with the general population. *The Lancet HIV*, 1(1), e32-e40.
21. Samson-Akpan, P. E., Ojong, I. N., Ella, R., & Edet, O. B. (2013). Quality of life of people living with HIV/AIDS in Cross River, Nigeria. *International Journal of Medicine and Biomedical Research*, 2(2), 207-212.
22. Lifson, A. R., Grandits, G. A., Gardner, E. M., Wolff, M. J., Pulik, P., Routy, J. P., & INSIGHT START Study Group. (2015). Quality of life assessment among HIV-positive persons entering the INSIGHT Strategic Timing of AntiRetroviral Treatment (START) trial. *HIV Medicine*, 16(1), 88-96.
23. Jadhav, P. S., Laad, P. S., & Chaturvedi, R. M. (2017). Quality of Life Factors Affecting Quality of Life in People Living with HIV / AIDS in An Urban Area. *International Journal of Community Medicine and Public Health*, 4(8), 3031-3036.
24. Anand, D., Puri, S., & Mathew, M. (2012). Assessment of quality of life of HIV-positive people receiving art: An Indian perspective. *Indian Journal of Community Medicine*, 37(3), 165-169.
25. Bello, S. I., & Bello, I. K. (2013). Quality of life of HIV/AIDS patients in a secondary health care facility, Ilorin, Nigeria. *Proceedings (Baylor University. Medical Center)*, 26(2), 116-119.
26. dos Santos, E. C. M., França Júnior, I., & Lopes, F. (2007). Quality of life of people living with HIV/AIDS in São Paulo, Brazil. *Revista de Saúde Pública*, 41(Supl 2), 64-71.
27. Bakiono, F., Ouédraogo, L., Sanou, M., Samadoulougou, S., Wendpouriré, P., Guiguemdé, L., & Simporé, J. (2014). Quality of life with HIV: a cross-section study in Ouagadougou, Burkina Faso. *SpringerPlus*, 3(1), 372.
28. Motilewa, O. O., Ekanem, U. S., Onayade, A., & Sule, S. S. (2015). A Comparative Study of Health Related-Quality of Life Among HIV Patients on Pre-HAART and HAART in Uyo South-South Nigeria. *Journal of Antivirals & Antiretrovirals*, 7(3), 60-68.
29. Shan, D., Ge, Z., Ming, S., Wang, L., Sante, M., & Wang, X. (2011). Quality of life and related factors among HIV-positive spouses from serodiscordant couples under antiretroviral therapy in Henan Province, China. *PLoS ONE*, 6(7), e21839.
30. Nojomi, M., Anbary, K., & Ranjbar, M. (2008). Health-related quality of life in patients with HIV/AIDS. *Archives of Iranian Medicine*, 11(6), 608-612.
31. Monteiro, F., Canavarro, M. C., & Pereira, M. (2016). Factors associated with quality of life in middle-aged and older patients living with HIV. *AIDS Care*, 28(1), 92-98.
32. Belak Kovacević, S., Vurusić, T., Duvancić, K., & Macek, M. (2006). Quality of life of HIV-infected persons in Croatia. *Collegium Antropologicum*, 30(1), 79-84.
33. Dalmida, S. G., Koenig, H. G., Holstad, M. M., & Thomas, T. L. (2015). Religious and Psychosocial Covariates of Health-Related Quality of Life in People Living with HIV/AIDS. *HIV/AIDS Research and Treatment Open Journal*, 1(1), 1-7.
34. Thulasingham, M., Premarajan, K. C., & Hamide, A. (2011). Quality of life and its determinants in people living with human immunodeficiency virus infection in puducherry, India. *Indian Journal of Community Medicine*, 36(3), 203-207.
35. Imam, M. H., Karim, M. R., Ferdous, C., & Akhter, S. (2011). Health related quality of life among the people living with HIV. *Bangladesh Medical Research Council Bulletin*, 37(1), 1-6.
36. Mutabazi-Mwesigire, D., Katamba, A., Martin, F., Seeley, J., & Wu, A. W. (2015). Factors That Affect Quality of Life among People Living with HIV Attending an Urban Clinic in Uganda: A Cohort Study. *PLoS ONE*, 10(5), e0126810.
37. Oliveira, F. B. M., Moura, M. E. B., de Araújo, T. M. E., & Andrade, E. M. L. R. (2015). Quality of life and associated factors in people living with HIV/AIDS. *Acta Paulista de Enfermagem*, 28(6), 510-516.
38. Maimaiti, R., Yuexin, Z., Kejun, P., Wubili, M., Lalanne, C., & Abudureyimu, K. (2017). Assessment of Health-Related Quality of Life among People Living with HIV in Xinjiang, West China. *Journal of the International Association of Providers of AIDS Care (JIAPAC)*, 16(6), 588-594.
39. Mahajan, A. P., Sayles, J. N., Patel, V. A., Remien, R. H., Sawires, S. R., Ortiz, D., & Rabkin, M. (2008). Stigma in the HIV/AIDS epidemic: A review of the literature and recommendations for the way forward. *AIDS*, 22(Suppl 2), S67-S79.
40. Webel, A. R., Longenecker, C. T., Gripshover, B., Hanson, J. E., Schmotzer, B. J., Salata, R. A., & стержень, М. В. (2014). Age, stress, and isolation in older adults living with HIV. *AIDS Care*, 26(4), 523-531.
41. Li, Z.-w., Jin, G., Yuan, J., Ma, X.-x., & Meng, X.-l. (2017). Impact Factors of Quality of Life among HIV/AIDS Patients after Antiretroviral Therapy: A Cohort Study Systematic Review. In *2nd International Conference on Information Technology and Management Engineering* (pp. 387-393).
42. Rai, Y., Tanusree, D., & Anit, K. G. (2010). Quality of life of HIV-infected people across different stages of infection. *Journal of Happiness Studies*, 12(1), 61-69.
43. Wig, N., Lekshmi, R., Pal, H., Ahuja, V., Mittal, C. M., Amte, A., & Sreenivas, V. (2006). The impact of HIV/AIDS on the quality of life: a cross sectional study in north India. *Indian Journal of Medical Sciences*, 60(1), 3-12.
44. Tran, B. X., & Nguyen, N. P. T. (2012). Patient Satisfaction with HIV/AIDS Care and Treatment in the Decentralization of Services Delivery in Vietnam. *PLoS ONE*, 7(9), e46680.
45. Yakob, B., & Ncama, B. P. (2016). Client satisfaction: correlates and implications for improving HIV/AIDS treatment and care services in southern Ethiopia. *International Health*, 8(4), 292-298.

46. World Health Organization. (2017). HIV/AIDS Fact sheet [Internet]. Geneva: World Health Organization Media Centre. Retrieved from <http://www.who.int/mediacentre/fact-sheets/fs360/en/>
47. The Joint United Nations Programme on HIV/AIDS (UNAIDS). (2017). Fact sheet - Latest statistics on the status of the AIDS epidemic-GLOBAL HIV STATISTICS [Internet]. Geneva. Retrieved from <http://www.unaids.org/en/resources/fact-sheet>
48. Basavaraj, K. H., Navya, M. A., & Rashmi, R. (2010). Quality of life in HIV/AIDS. *Indian Journal of Sexually Transmitted Diseases and AIDS*, 31(2), 75-80.
49. Jia, H., Uphold, C. R., Wu, S., Reid, K., Findley, K. (2004). Health-related quality of life among men with HIV infection: effects of social support, coping, and depression. *AIDS Patient Care and STDs*, 18(10), 594-603.
50. World Health Organization. (1948). Constitution of the World Health Organization. Geneva.
51. Olusina, A. K., & Ohaeri, J. U. (2003). Subjective quality of life of recently discharged Nigerian psychiatric patients. *Social Psychiatry and Psychiatric Epidemiology*, 38(12), 707-714.
52. World Health Organization. (1996). WHOQOL-BREF. Introduction, Administration, Scoring and Generic version of the Assessment. Field Trial Version [Internet]. Geneva. Retrieved from http://www.who.int/mental_health/media/en/76.pdf
53. World Health Organization. (2002). WHOQOL-HIV BREF. Geneva.
54. Mills, E. J., Bakanda, C., Birungi, J., Chan, K., Ford, N., Cooper, C. L., & Nachega, J. B. (2011). Life Expectancy of Persons Receiving Combination Antiretroviral Therapy in Low-Income Countries: A Cohort Analysis from Uganda. *Annals of Internal Medicine*, 155(4), 209-216.
55. Poupard, M., Ngom Gueye, N. F., Thiam, D., Ndiaye, B., Girard, P. M., & Laurent-Lewin, V. (2007). Quality of life and depression among HIV-infected patients receiving efavirenz- or protease inhibitor-based therapy in Senegal. *HIV Medicine*, 8(2), 92-95.
56. Stangl, A. L., Wamai, N., Mermin, J., Awor, A. C., & Bunnell, R. E. (2007). Trends and predictors of quality of life among HIV-infected adults taking highly active antiretroviral therapy in rural Uganda. *AIDS Care*, 19(5),
57. Degroote, S., Vogelaers, D., & Vandijck, D. M. (2014). What determines health-related quality of life among people living with HIV: an updated review of the literature. *Archives of Public Health*, 72(1), 40.
58. Merkouris, A., Andreadou, A., Athini, E., Hatzimbalasi, M., Rovithis, M., & Papastavrou, E. (2013). Assessment of patient satisfaction in public hospitals in Cyprus: a descriptive study. *Health Science Journal*, 7(1), 28-40.
59. Deeks, S. G., Lewin, S. R., & Havlir, D. V. (2013). The end of AIDS: HIV infection as a chronic disease. *The Lancet*, 382(9903), 1525-1533.
60. Centers for Disease Control and Prevention. (2016). HIV Among People Aged 50 and Over. U.S. Department of Health and Human Services.
61. Enimil, A., Nugent, N., Amoah, C., Norman, B., Antwi, S., Appiah, L., & Renner, L. (2016). Quality of life among Ghanaian adolescents living with perinatally acquired HIV: a mixed methods study. *AIDS Care*, 28(4), 460-464.
62. Louw, J., Peltzer, K., Naidoo, P., Matseke, G., Mchunu, G., & Zuma, K. (2012). Quality of life among tuberculosis (TB), TB retreatment and/or TB-HIV co-infected primary public health care patients in three districts in South Africa. *Health and Quality of Life Outcomes*, 10(1), 77.
63. Katon, W. J. (2011). Epidemiology and treatment of depression in patients with chronic medical illness. *Dialogues in Clinical Neuroscience*, 13(1), 7-23.
64. Jelsma, J., MacLean, E., Hughes, J., Tinise, X., & Darder, M. (2005). An investigation into the health-related quality of life of individuals living with HIV who are receiving HAART. *AIDS Care*, 17(5), 579-588.
65. Wu, A. W. (2000). Quality of life assessment comes of age in the era of highly active antiretroviral therapy. *AIDS*, 14(14), 2449-2451.
66. Briongos Figuero, L. S., Bachiller Luque, P., Palacios Martín, T., González Sagrado, M., & Eiros Bouza, J. M. (2011). Assessment of factors influencing health-related quality of life in HIV-infected patients. *HIV Medicine*, 12(1), 22-30.
67. Hays, R. D., Cunningham, W. E., Sherbourne, C. D., Wilson, I. B., Wu, A. W., Cleary, P. D., & Shapiro, M. F. (2000). Health-related quality of life in patients with human immunodeficiency virus infection in the United States: results from the HIV Cost and Services Utilization Study. *The American Journal of Medicine*, 108(9), 714-722.
68. Tsevat, J., Leonard, A. C., Szaflarski, M., Sherman, S. N., Cotton, S., точки, Д., & Giordano, T. P. (2009). Change in quality of life after being diagnosed with HIV: a multicenter longitudinal study. *AIDS Patient Care and STDs*, 23(11), 931-937.
69. UNAIDS. (2000). National AIDS Programmes: A guide to monitoring and evaluation. Retrieved from http://www.who.int/hiv/pub/epidemiology/en/JC427-Mon_Ev-Full_en.pdf?ua=1
70. Drummond, M. F., Sculpher, M. J., Claxton, K., Stoddart, G. L., & Torrance, G. W. (2015). *Methods for the economic evaluation of health care programmes* (4th ed.). Oxford University Press.
71. Nakagawa, F., Lodwick, R. K., Smith, C. J., Smith, R., Cambiano, V., Lundgren, J. D., & группа сотрудничества D:A:D. (2012). Projected life expectancy of people with HIV according to timing of diagnosis. *AIDS*, 26(3), 335-343.
72. Ndubuka, N. O., Lim, H. J., Ehlers, V. J., & van der Wal, D. M. (2017). Health-related quality of life of patients on antiretroviral treatment in Botswana: A cross-sectional study. *Palliative and Supportive Care*, 15(2), 214-222.
73. Do, A. N., Rosenberg, E. S., Sullivan, P. S., Beer, L., Strine, T. W., & Hall, H. I. (2014). Excess burden of depression among HIV-infected persons receiving medical care in the United States: data from the medical monitoring project and the behavioral risk factor surveillance system. *PLoS One*, 9(3), e92842.
74. Duracinsky, M., Lalanne, C., Le Coeur, S., Herrmann, S., Berzins, B., چنس, C., & فورگ PROQOL-HIV. (2012). Psychometric validation of the PROQOL-HIV questionnaire, a new health-related quality of life instrument-specific to HIV disease. *Journal of Acquired Immune Deficiency Syndromes*, 59(5), 506-515.

75. Reis, R. K., Santos, C. B., & Gir, E. (2012). Quality of life among Brazilian women living with HIV/SIDA. *AIDS Care*, 24(5), 626-634.
76. da Silva, J., Bunn, K., Bertoni, R. F., Neves, O. A., & Traebert, J. (2013). Quality of life of people living with HIV. *AIDS Care*, 25(1), 71-76.
77. Catherine, W., & Hartmut, B. K. (2004). Employment status, level of income and quality of life in people living with HIV. Retrieved from <http://www.pulsus.com/cahr2004/abs401.htm>
78. Hipolito, R. L., & de Oliveira, D. C. (2016). Quality of Life of people living with HIV/AIDS: a cross-sectional study. *Brazilian Journal of Nursing*, 15(3), 575-578.
79. Mofolorunsho, K. C., Nwankwo, E., & Mofolorunsho, T. B. (2013). Socio-Economic Factors Influencing the Quality of Life of People Living with HIV/AIDS in Kogi, Nigeria. *Nature and Science*, 11(1), 33-39.
80. National Population Commission, Nigeria, & ICF Macro. (2008). *Nigeria Demographic and Health Survey 2008*. Abuja, Nigeria: National Population Commission and ICF Macro.
81. Ana, C. R., Leonor, L., Marina, P. G., & Eduardo, R. (2010). Relationship among psychopathological symptoms, treatment adherence and quality of life in HIV/AIDS infection. *Psicologia: Reflexão e Crítica*, 23(3), 420-429.
82. Adebajo, S. B., Bamgbala, A. O., & Oyediran, M. A. (2003). Attitudes of health care providers to persons living with HIV/AIDS in Lagos State, Nigeria. *African Journal of Reproductive Health*, 7(3), 103-112.
83. Thulasigam, M., Premarajan, K., & Hamide, A. (2011). Quality of life and its determinants in people living with human immunodeficiency virus infection in puducherry, India. *Indian Journal of Community Medicine*, 36(3), 203-207.
84. Mannheimer, S. B., Matts, J., Telzak, E., Chesney, M., Child, C., Wu, A. W., & Friedland, G. (2005). Quality of life in HIV-infected individuals receiving antiretroviral therapy is related to adherence. *AIDS Care*, 17(1), 10-22.
85. Willis, N., Mavhu, W., Wogrin, C., Mutsinze, A., & Kagee, A. (2018). Understanding the experience and manifestation of depression in adolescents living with HIV in Harare, Zimbabwe. *PLoS One*, 13(1), e0190423.
86. Chesney, M. A., & Smith, A. W. (1999). Critical delays in HIV testing and care. The potential role of stigma. *American Behavioral Scientist*, 42(7), 1162-1174.
87. Wingood, G. M., DiClemente, R. J., Mikhail, I., Hubbard McCree, D., Davies, S. L., Harrington, K. F., & думал, H. (2008). HIV discrimination and the health of women living with HIV. *Women & Health*, 46(2), 99-112.
88. Ma, L., Xu, P., Lin, H., Ju, L., & Lv, F. (2015). Quality of Life of People Living with HIV/AIDS: A Cross-Sectional Study in Zhejiang Province, China. *PloS One*, 10(8), e0135705.
89. Mutabazi-Mwesigire, D., Seeley, J., Martin, F., & Katamba, A. (2014). Perceptions of quality of life among Ugandan patients living with HIV: a qualitative study. *BMC Public Health*, 14(1), 343.
90. Wilson, I. B., & Cleary, P. D. (1995). Linking clinical variables with health-related quality of life: A conceptual model of patient outcomes. *JAMA*, 273(1), 59-65.
91. Bajunirwe, F., Bangsberg, D. R., & Sethi, A. K. (2013). Alcohol use and HIV serostatus of partner predict high-risk sexual behavior among patients receiving antiretroviral therapy in South Western Uganda. *BMC Public Health*, 13(1), 430.
92. Drewes, J., Gusy, B., & von Rüden, U. (2013). More than 20 years of research into the quality of life of people with HIV and AIDS—a descriptive review of study characteristics and methodological approaches of published empirical studies. *Journal of the International Association of Providers of AIDS Care (JIAPAC)*, 12(1), 18-22.
93. Herrmann, S., McKinnon, E., Hyland, N. B., Lalanne, C., Mallal, S., & Stewart, A. (2013). HIV-related stigma and physical symptoms have a persistent influence on health-related quality of life in Australians with HIV infection. *Health and Quality of Life Outcomes*, 11(1), 56.
94. Karkashadze, E., Gates, M. A., Chkhartishvili, N., DeHovitz, J., & Tsertsvadze, T. (2017). Assessment of quality of life in people living with HIV in Georgia. *International Journal of STD & AIDS*, 28(7), 672-678.
95. Peltzer, K., & Phaswana-Mafuya, N. (2008). Health-related quality of life in a sample of HIV-infected South Africans. *African Journal of AIDS Research*, 17(2), 209-218.
96. Yan, Z., Wan, D., & Li, L. (2011). Patient satisfaction in two Chinese provinces: rural and urban differences. *International Journal for Quality in Health Care*, 23(4), 384-389.
97. Chimbindi, N., Bärnighausen, T., & Newell, M.-L. (2014). Patient satisfaction with HIV and TB treatment in a public programme in rural KwaZulu-Natal: evidence from patient-exit interviews. *BMC Health Services Research*, 14(1), 32.
98. Vo, B. N., Cohen, C. R., Smith, R. M., Bukusi, E. A., Onono, M. A., Turan, J. M., & Edmonds, A. (2012). Patient satisfaction with integrated HIV and antenatal care services in rural Kenya. *AIDS Care*, 24(11), 1442-1447.
99. Belay, M. (2013). HIV/AIDS Patients' satisfaction on ART laboratory service in selected governmental hospitals, Sidamma Zone, southern Ethiopia. *Science Journal of Public Health*, 1(3), 85-90.
100. Okoye, M. O., Ukwe, V. C., Okoye, T. C., Adibe, M. O., & Ekwunife, O. I. (2014). Satisfaction of HIV patients with pharmaceutical services in South Eastern Nigerian hospitals. *International Journal of Clinical Pharmacy*, 36(1), 1-8.
101. Chimbindi, N., Bärnighausen, T., & Newell, M.-L. (2014). Patient satisfaction with HIV and TB treatment in a public programme in rural KwaZulu-Natal: evidence from patient-exit interviews. *BMC Health Services Research*, 14(1), 32.
102. Ndou, T. V., Maputle, S. M., & Risenga, P. R. (2016). HIV-positive patients' perceptions of care received at a selected antiretroviral therapy clinic in Vhembe district, South Africa. *African Journal of Primary Health Care and Family Medicine*, 8(1), a926.
103. Nabbuye-Sekandi, J., Makumbi, F. E., Kasangaki, A., Kizza, I. B., & Tugumisirize, J. (2011). Patient satisfaction with services in outpatient clinics at Mulago hospital, Uganda. *International Journal for Quality in Health Care*, 23(5), 516-523.
104. Belay, M., Abrar, S., Bekele, D., Daka, D., Derbe, M., & et al. (2013). HIV/AIDS Patients' satisfaction on ART laboratory service in selected governmental hospitals, Sidamma Zone, southern Ethiopia. *Science Journal of Public Health*, 1(3), 85-90.

105. Wung Buh, A., Fon Peter, N., & Atashili, J. (2016). Clients' satisfaction with HIV treatment services in Bamenda, Cameroon: a crosssectional study. *BMC Health Services Research*, 16(1), 280.
106. Oche, M., Raji, M., Kaoje, A., Gana, G., Ango, J., & Geidam, Y. A. (2013). Clients' satisfaction with anti-retroviral therapy services in a tertiary hospital in Sokoto, Nigeria. *Journal of AIDS and HIV Research*, 5(11), 328-333.
107. Nwabueze, S. A., Adogu, P. O., Ilika, A. L., Asuzu, M. C., & Adinma, E. D. (2011). Perception of quality of care in HIV/AIDS programmes among patients in a tertiary health care facility in Anambra State. *Nigerian Journal of Medicine*, 20(1), 144-150.
108. World Health Organization. (2016). Consolidated Guidelines on the Use of Antiretroviral Drugs for Treating and Preventing HIV Infection: Recommendations for a Public Health Approach (2nd ed.).
109. Wilson, I. B., & Cleary, P. D. (1995). Linking clinical variables with health-related quality of life: A Conceptual model of patient outcomes. *JAMA*, 273(1), 59-65.
110. Ferrans, C. E., Zerwic, J. J., Wilbur, J. E., & Larson, J. L. (2005). Conceptual model of health-related quality of life. *Journal of Nursing Scholarship*, 37(4), 336-342.
111. Walker, L. O., & Avant, K. C. (2005). *Strategies for theory construction in nursing* (4th ed.). Prentice Hall.
112. Bakas, T., McLennon, S. M., Carpenter, J. S., Buelow, J. M., Otte, J. L., & Yeager, K. A. (2012). Systematic review of health-related quality of life models. *Health and Quality of Life Outcomes*, 10(1), 134.
113. National Population Commission, Nigeria, & ICF International. (2014). *Nigeria Demographic and Health Survey 2013*. Abuja, Nigeria: National Population Commission and ICF International.