

Exploring Barriers and Perceptions of Rural Women Towards Cervical Cancer Screening Among Reproductive Age Women in Sadi Chanka District, Oromia, West Ethiopia: A Qualitative Study

Ararso Hordofa Guye*, Dame Banti Shambi, Tadesse Nigussie, Elisabeth Lagesse, Negasa Shuma Dureso, Efa Bayissa Kanea, and Kasim Hansa

Department of Public Health, College of Medicine and Health Sciences, Salale University, Fitcha, Ethiopia

***Corresponding author:** Ararso Hordofa Guye, Department of Public Health, College of Medicine and Health Sciences, Salale University, Fitcha, Ethiopia.

Submitted: 01 September 2023 **Accepted:** 08 September 2023 **Published:** 13 September 2023

Citation: Ararso Hordofa Guye, Dame Banti Shambi, Tadesse Nigussie, Elisabeth Lagesse, Negasa Shuma Dureso, et al. (2023) Exploring Barriers and Perceptions of Rural Women Towards Cervical Cancer Screening Among Reproductive Age Women in Sadi Chanka District, Oromia, West Ethiopia: A Qualitative Study. *Ethiopia. Sci Set J Cancer Res* 2(1), 01-05.

Abstract

Background: Cervical cancer is the fourth most common cancer affecting women's reproductive organs and challenges faced by women. Cervical cancer screening is a public health intervention used on a population at risk, or target population to diagnose a disease. Effective screening programs reduce morbidity and mortality due to this cancer. The study aimed to explore barriers and perceptions of rural women towards cervical cancer screening among reproductive age women in Sadi Chanka district, Oromia, West Ethiopia.

Methods: A qualitative study was conducted in Sadi Chanka district through in-depth interviews of 28 study participants from November 1 to December 20, 2022. Data were collected using semi-structured questionnaires and captured using an audio tape recorder and field note-taking. A homogeneous purposive sampling technique was used to select representative study participants. Participants' interviews were transcribed and translated according to the participants verbatim from the local language (Afan-Oromo) to the English language by language experts in both languages. The transcribed data were entered into Microsoft Word and analyzed using thematic analysis. Data were presented in narrative forms using the respondent's own verbatim as an illustration.

Results: A total of twenty-eight study participants were involved in this study. Seven thematic areas were emerged from interviews of the respondents through thematic analysis of the data: The identified barriers and perceptions of the respondents towards cervical screening were categorized into the following thematic areas are; Misinformation and awareness related factors, sign and symptoms of cervical cancer, gender preferred for screening service, side effects of procedures, availability and affordability of the service, and prevention and treatment outcome of the disease were the key identified barriers and perceptions of cervical cancer screening under these themes.

Conclusion: The finding explored and revealed that, misinformation, lack of awareness about cervical cancer screening services, lack of cervical screening services at all health facilities, risk factors of cervical cancer, sign and symptoms of cervical cancer, gender preferred for screening service, side effects of procedures, availability and affordability of the service, and prevention and treatment outcome of the disease were the key barriers and perceptions of cervical screening among rural reproductive age women. Thus, the district health office should work on cervical cancer screening services to improve rural women's barriers and perceptions of cervical cancer service screening, efforts should be focused on enhancing awareness and related factors.

Keywords: Cervical Cancer, Cervical Cancer Screening, Barriers, Perception, Exploring, Ethiopia

Abbreviations

CCS: Cervical Cancer Screening
HPV: Human Papilloma Virus
VIA: Visual Inspection of the Cervix with Acetic Acid
IDIs: In-Depth Interviews.

Introduction

Cervical cancer is the most common cancer affecting women's reproductive organs and also the fourth leading cause of death from cancer among women worldwide [1]. There are two main types of cervical cancer, squamous cell carcinomas which is the

most common type, and accounts for 90 percent of cases, and the second one is adenocarcinomas are cancers that develop from the glandular cell [2]. The primary cause of cervical cancer is human papillomavirus (HPV), HPV 16 and 18 were among the most common and prevalent types, responsible for approximately 70% of all cervical cancer cases and prevented through immunization with the HPV vaccine [3].

Cervical cancer screening (CCS) is a public health intervention used on populations at risk, or eligible populations to identify cervical abnormalities in an asymptomatic stage [4]. Effective CCS is reducing morbidity and mortality due to this cancer [1]. The Ethiopia Federal Minister of Health (FMoH) recommends that women begin screening for cervical cancer at the age of 30 and continue every five years by using visual inspection with acetic acid (VIA) [5].

Globally, cervical cancer kills an estimated 311,000 women and 570,000 new cases are reported in 2018 [6]. The burden of the disease in Low-middle income countries shared about 85% of the 520,000 morbidity and 87% of the 231,000-mortality due to cervical cancer every year in 2012, and in central and eastern Africa, 266,000 women death occur due to this cancer [7]. There are several risk factors for cervical cancer, sexual intercourse at an early age, multiple sexual partners, tobacco smoking behavior, long-term use of oral contraceptive pills, immune system compromise, chlamydia infection, HPV, and low socio-economic status [8]. In Ethiopia, nearly 20.9 million women were at risk of developing cervical cancer. Annually, 4,648 and 3,235 new cases and death occurred respectively. The prevalence of cervical cancer incidence was 13.4%, and much of this cancer occurred among women aged 15 to 44 years [1]. The coverage of CCS service availability is very low in Ethiopia, only nine percent of health facilities offered screening services and less than one percent of women undergo screening in the country [9, 10]. To improve the perceptions of women on cervical cancer and its screening, the Ethiopian government implements a national cancer control plan. But, still, the perceived status of women is not changed [11].

Some studies had been conducted in Ethiopia indicating barriers and perceptions of rural women towards cervical cancer screening as well as the level of CCS among reproductive age women [12, 13]. However, most of the studies are focused on urban women. But, this study focused on rural women. The findings from this study may provide some information on the barriers and perceptions of rural women toward cervical cancer screening. The study aimed to explore barriers and perceptions of rural women towards cervical cancer screening in Sadi Chanka District, Oromia, West Ethiopia.

Methods and Materials

Study Design, Period and Area

A qualitative phenomenological study design was conducted using in-depth interviews (IDIs) to explore barriers and perceptions of rural women's on cervical cancer screening among reproductive age women in Sadi Chanka district, Oromia, west Ethiopia from November 1 to December 20, 2022. Sadi Chanka district is located 588 Km from Addis Ababa and 66 Km from Dambi Dollo, the Capital city of Kellem Wollaga Zone. According to the Central Statistical Agency population projections of 2014, the total estimated population was 66,320. Of these 32,497 were women. According to updated data from the family folder of

health extension workers, there were 11,969 rural women aged 30-49 years in the district. The district has 15 administrative kebeles, 13 rural kebeles, and two urban kebeles. In the district, there are two governmental health centers and 15 health posts. From these health facilities, the health centers provide cervical cancer screening services by using VIA

Study Population

The study population was purposively selected rural women aged 30-49 years old living in eligible households from randomly selected rural kebeles in the district.

Eligibility Criteria

All rural women of aged 30-49 years, in selected kebele of Sadi Chanka district, lived for at least 6 months were included in the study. Rural women aged 30-49 years who were unable to respond and critically ill women were excluded from the study.

Sample Size and Sampling Technique

Twenty eight rural women in reproductive age women living in Sadi Chanka district were included in the study. The study setting has 13 administrative rural kebeles and the study was carried out in the five selected rural kebeles of the district. Five rural kebeles were randomly selected by lottery method from a total of fifteen rural kebeles in the district. A homogenous purposive sampling technique was used for selecting study participants from the rural kebeles to get the women who would be willing to share the barriers and their perceptions of cervical cancer screening. The included participants were reproductive age women in each of the five selected rural kebeles and the sample size was determined by the saturation idea of the participants IDIs.

Data Collection Tools and Procedures

Data were collected by using a semi-structured interview guide for IDIs for rural reproductive age women with flexible probing techniques except for the background data. An interview guide was developed by reviewing different kinds of literature [12, 13]. The data were collected through face-to-face interviews and the IDIs were conducted by the three master's holder health professionals who are fluent in the local language. Data were collected by using interview guide, field note-taking and tape-recording for each participant. Data were collected by using interview guide and an audio tape recorder and field note-taking for each participant, and the interview time ranged from 30 minutes to 60 minutes based on the saturation of the participants' ideas. The IDIs were conducted in a place convenient for the respondents are quiet and confidential setting. The transcription and translation of the data were done from an audio tape recorder for each participant's word-by-word according to their verbatim responses. Four master's holder health professionals who are fluent in local languages conducted in-depth interviews with rural reproductive age women. The transcription and translation of data were done from an audio tape recorder for each respondent word by word according to their verbatim responses.

Data Quality Assurance

The semi-structured interview guide was prepared in English and translated into the local language (Afan-Oromo) and then back-translated into English to ensure consistency. Finally, the Afan Oromo version of interview guide was used for data collection. Before actual data collection, the data collectors received two days of training on the objective of the study, data gathering methods, and ethical considerations. The interview guides were

evaluated by qualitative research experts. During interviews, probing techniques were used and the interviewers used tape-record and not-taking. The IDIs took place at the respondent's houses in a private and confidential setting. Another healthcare practitioner double-checked the content validity of the tool's. Before being used for actual data collection, the instruments were pretested among rural reproductive age women that were similar to those found in another district. By choosing and maintaining a peaceful environment for interviews and audio recording with chosen study participants, the quality of the data was ensured. At the time of the interviews, each participant's thorough handwritten field notes were taken. The participants were urged to speak up, express themselves openly, and share their own experiences with cervical cancer screening. All audio recordings of the interviews were translated into the participants' native language, Afan Oromo, before being used.

Data Processing and Analysis

The collected data were first transcribed according to the verbatim of the respondents into local language Afaan Oromo, by which the interviews were conducted. Next, the interviewed data were translated and transcribed into English by senior language experts. Then, a final edition of the code was developed, and the categories and themes were constructed. Data analysis was started during data collection and then transcribed and translated to English language by replaying the recorded audio tape. Informed consent, transcribed data, and audio record files were stored in a private place. Data were coded, categorized, and analyzed by using thematic analysis manually and finally, the result was presented by narrative forms to describe the barriers and perceptions of cervical cancer screening

Results

Socio-demographic Characteristics of Participants

A total of 28 rural reproductive age women participated in this study. Their ages ranged between 33 to 48 years. Regarding the marital status of the respondents, 25(89.2%) were marrieds. The majority of the respondents, 18 (64.2 %) had not attended formal education. Of the total respondents, 21(75%) were Protestant Christians. Seven thematic areas that emerged from the IDIs were captured and organized as reported below.

Theme: Misinformation and Awareness Related Factors

The finding from the in-depth interviews of rural reproductive age women indicates that misinformation and lack of awareness about cervical cancer screening were the factor for barriers and perceptions of rural reproductive age women. The 34 years rural women stated that: *".... including me, most of the rural women did not give attention for cervical cancer screening due to lack of awareness and not knowing the benefit of cervical cancer screening as well...and some of the women have misinformation about cervical cancer screening."*

The 38 years old women said that: *"....due to we are living in rural area we are not getting adequate information on cervical cancer screening. The health extension workers are also not providing cervical cancer related health education for us and do not give enough explanation in detail to the rural women."*

Theme: Risk Factors of Cervical Cancer

Concerning the risk factors of cervical cancer thematic area, the evidence from IDIs of the respondents indicated that lack of awareness of cervical cancer causes, susceptibility, and who

is at risk for the disease were the reasons for women not utilizing cervical cancer screening services. Most of the respondents perceived that they were not susceptible to this cancer, so the screening service is not required.

One of the key informant interviews raised the idea that: *"I am not feared of this cancer because I have no chance of developing cervical cancer..... and I have no detail understanding on cervical cancer"* (44 years-old women, IDIs participant).

Theme: Signs and Symptoms of the Cervical Cancer

Sign and symptoms of cervical cancer was the other thematic area identified from the IDIs of the respondents. The majority of respondents raised the absence of signs and symptoms of the disease are the reason for not being screened for cervical cancer.

One of the key informant interviews said that: *"I think the main cause for many women is not screened for cervical cancer is due to the natural history of cancer. I mean, at that period they were healthy and had no signs, symptoms, or complications of the disease. Our women are more concerned if there are detectable signs and symptoms of the disease."* (48 years-old woman, IDIs participant).

Theme: Gender Preferring Service

The participants report different perceptions about cervical cancer screening services. From these, the fear of health care providers was identified as a factor for women not utilizing cervical cancer screening. A 36 years-old woman mentioned: *"Before five months, I was screened for cancer of the cervix by a male Doctor and I was ashamed when he asked me to remove my clothes...."* (36 years-old woman, IDIs participant)

Theme: Side Effects of Procedures

The evidence from the ideas of women showed that misinformation and rumors about the procedures of cervical cancer screening affect rural women's perception of cervical cancer and its screening. Most of the study participants interviewed raised the main reason for women not utilizing screening services was due to fear of the side effects of screening like vaginal bleeding, pain during and after the screening, and infertility.

The 42 years-old rural women reported that *"I heard some women told me that the screening procedures may cause excessive vaginal bleeding, painful..."* Another key informant interview raised the idea that: *"Cervical cancer screening procedures could lead to health issues. Such as infertility...."* (38 years-old women, IDIs participant).

Theme: Availability and Affordability of the Service

The finding from this study also indicated that rural women not having sufficient awareness about the availability and affordability of the service was the reason for women not having a high perception of cervical cancer and its screening.

One of the key informant interviews said that: *"I don't know that the cervical cancer screening service is given freely in the health facility, I think the cervical cancer screening services are given by paying the costs....."* (33 years-old women, IDIs participant).

"I heard that the payment for cervical cancer screening service is very expensive." (40 years-old women, IDIs participant). Another key informant interviews raised the idea: *"....two years*

ago, I was treated for vaginal discharge and abdominal pain at the health center; then I was wanted to screen for cervical cancer disease after counseling by Doctors, but I don't know the availability of the service at that health facility.” (45 years-old women, IDIs participant)

Theme: Prevention and Treatment Outcome of the Disease

The sixth theme is perceptions about the prevention and treatment outcome of the disease. Most of the women perceived that cervical cancer is not curable and cannot be prevented by screening.

One respondent said that: *“I think screening for cervical cancer disease is not essential for healthy women and women of this age group...”* (39 years-old woman, IDIs participant)

Another 46 years old woman stated that *“...as women most of a time we believe that cervical cancer screening is used for women who are ill or who are approaching menopause.”*

According to the barriers and perceptions of rural women, cervical cancer is not prevented by treatment and screening. They linked it with religion and belief and most of women prefer to pray to God instead of screening for cervical cancer and treating the disease. They also think and believed that cervical cancer treatment outcome is useless even if treated.

“...I choose to die by the disease rather than utilizing the cervical cancer screening and being told having cancer problem due to some of health care providers not keep the confidentiality of the patients...and their treatment is not satisfactory” (35 years-old women, IDIs participant)

Discussion

This study aimed to explore barriers and perceptions of rural reproductive age women towards cervical cancer screening in Sadi Chanka district, Oromia, west Ethiopia. This study identified the key rural women's perceptions of cervical cancer screening and the identified barriers and perceptions of cervical cancer screening among rural women were related to the following seven themes: misinformation and awareness related factors, risk factors of cervical cancer, sign and symptoms of cervical cancer, gender preferred for screening service, side effects of procedures, availability and affordability of the service, and prevention and treatment outcome of the disease.

The result of this study showed that the majority of rural women had a many barriers and perceptions of cervical cancer screening with the risk factors of cervical cancer. Some rural women were unaware of the cause of cervical cancer or who was at risk for this cancer. This result agrees with the results of studies done in the Wolaita zone, Addis Ababa, Debre Berhan, and Malawi [12-16]. This might be due to lower awareness about the risk factors of this cancer.

Regarding the barriers and perceptions of the signs and symptoms of cervical cancer, some of the rural women were a low perception of the disease's signs and symptoms. This is comparable to a study done in the Wolaita zone and Ambo town [12, 17]. This could be due to low attention to health education about cervical cancer and its screening given by media and health professionals.

According to this study, some participants wanted to be screened by female healthcare providers because most women fear being

unclothed in front of male healthcare providers. This is similar to a study conducted in Jimma, Kenya, Nigeria and Cameroon [18-21]. This could be because religions and cultures have negatively influenced women health-seeking behaviors, practices, actions, beliefs, gender roles, and self-confidence.

The study participants had negatively perceived cervical cancer screening procedures. Most rural women think screening procedures may cause infertility, abnormal vaginal bleeding, and pain during the examination. This finding is also consistent with finding in Malawi and Kenya [16, 19]. This association can be explained by women not having enough information and misunderstanding about the producers of cervical cancer screening.

Concerning the availability and affordability of the service, many women perceived cervical cancer screening service does not exist in the study area and that the screening is not a free service. This is also in line with the study done in Cameroon [13, 20]. This may happen due to a lack of access to information about existing and payment of screening services by health care providers. More women perceived that cervical cancer is not a curable disease and could not be prevented by treatment or screening. This is supported by the study conducted in the Wolaita zone and east Gojjam zone [12, 21]. This implies a lack of knowledge about the method of prevention and outcome of the treatments for cervical cancer [22].

Limitations of the Study

This study has encountered some limitations such as most of the discussion part was compared with the mixed report due to a limited number of qualitative studies done on the barriers and perceptions of rural women about cervical cancer screening. This study has its limitations as that of qualitative finding nature and the study samples were determined by the saturation idea of the respondents and it might not be representative of the general diverse population, and the results cannot be generalized to other settings

Conclusion

The study showed that rural reproductive age women in the study area had the following barriers and perceptions cervical cancer screening; misinformation and awareness related factors, the risk factors of cervical cancer, sign and symptoms of cervical cancer, gender preference for screening service, side effects of procedures, availability and affordability of the service, and prevention and treatment outcome of the diseases were the key barriers and perceptions of women for cervical cancer screening. Thus, the district health office should work on cervical cancer screening services to improve rural women's barriers and perceptions of cervical cancer service screening, efforts should be focused on enhancing awareness and related factors with cervical cancer screening.

Ethical Approval

The ethical approval for the study was obtained from the Salale University College of Health Science's ethical review committee before starting data collection. Letter of permission was received from all selected kebeles (villages) and written informed consent was obtained from all study participants after an explanation of the purpose and procedure of the study.

Authors' Contributions

AHG, DBS and TN participated in conceptualization, data curation, formal analysis, investigation, funding acquisition, meth-

odology, project administration, resources, software, supervision, and validation, visualization, writing original draft, writing review & editing of the manuscript. NSD, EL, EBK, and KH were participated in the methodology, investigation, resources, data curation, supervision, validation, review, and editing of the manuscript

Data Sharing Statement

Data produced or analyzed during this study are included in this published paper. The datasets can be available from the corresponding author based on reasonable request.

Consent for publication

Not applicable

Funding

The financial fund for this research work was covered by Salale University

Disclosure

The authors declared that they have no conflicts of interest for this work.

Acknowledgement

We would like to express our deepest gratitude and thanks to Salale University for providing us the opportunity to participate on research work activities and financial support. Finally, our sincere heartfelt thanks go to the data collectors, supervisors, study participants, and the Sadi Chanka District Health Office for their kind supports during conducting of this study

References

1. Federal Ministry of Health of Ethiopia (FMOH) 92015) Guideline for Cervical Cancer Prevention and Control in Ethiopia.
2. American Cancer Society (2017) What is Cervical Cancer? J Gynecol Womens Heal 2: 1-9.
3. Pan American Health Organization (2018) Plan of action for cervical cancer prevention and control 2018-2030. 1-26.
4. World Health Organization (WHO) (2018) Cervical Cancer Guidance for Operation: Cervical cancer prevention, screening and treatment of pre-cancer, diagnosis and treatment of cancer Guidance for MSF Operations. Msf 1-9.
5. Federal Democratic Republic of Ethiopia Ministry of Health (2015) National guidelines for the management of sexually transmitted infections using syndromic approach. Fed Minist Heal.
6. World Health Organization (WHO) (2020) Global strategy to accelerate the elimination of cervical cancer as a public health problem 1-56.
7. World Health Organization (WHO) (2014) Guidelines for Comprehensive Cervical Cancer Control 2: 1-408.
8. American Cancer society (2020) Cervical Cancer Causes, Risk Factors, and Prevention Risk Factors for Cervical Cancer. 1-15.
9. Ethiopian Public Health Institute (2018) Services Availability and Readiness Assessment (SARA) Ethiopian Public Health Institute Ethiopia Service Availability and Readiness Assessment (SARA) 2018 Final Report. 1-119.
10. Gakidou E, Nordhagen S, Obaermeyer Z (2008) Coverage of Cervical Cancer Screening in 57 Countries: Low Average Levels and Large Inequalities. PLoS One 5: 863-868.
11. Federal Ministry of Health Ethiopia (2020) National cancer control plan 2016-2020.
12. Azeze GA, Damissie BW, Asseffa NA, Lake EA, Besha BB, et al. (2022) Communities' perceptions towards cervical cancer and its screening in Wolaita zone, southern Ethiopia: A qualitative study. PLoS One 17: 1-12.
13. Gizaw M, Ayele W, Getachew E, Getachew S, Addissie A, et al. (2019) Cervical cancer screening knowledge and barriers among women in Addis Ababa. PLoS One 7: 1-13.
14. Gonie A, Worku C, Assefa T, Bogale D, Girma A (2018) Acceptability and factors associated with post-partum IUCD use among women who gave birth at bale zone health facilities, Southeast-Ethiopia. Contracept Reprod Med 3: 1-8.
15. Mruts KB, Gebremariam TB (2018) Knowledge and Perception Towards Cervical Cancer among Female Debre Berhan University Students. Asian Pacific J Cancer Prev 19: 1771-1777.
16. Bula AK, Lee F, Chapola J, Mapanje C, Tsidya M, et al. (2022) Perceptions of cervical cancer and motivation for screening among women in Rural Lilongwe, Malawi: A qualitative study. PLoS One 1-13.
17. Natae SF, Nigatu DT, Negawo MK, mengesha WW (2020) Cervical Cancer Screening Uptake and Determinant Factors Among Women in Ambo Town: Community based Cross Sectional Study. Res Sq 1-17.
18. Nugusse T, Admassu B, Nigussie A (2019) Cervical cancer screening service utilization and associated factors among age-eligible women in Jimma town using health belief. BMC Womens Health 1-10.
19. Muiruri P, Njuguma E, Ilovi S, Mutai K, Kinuthia J, et al. (2017) Factors influencing cervical cancer screening in a Kenyan Health Facility: a mixed qualitative and quantitative study. ResearchGate.
20. Wilson AT, Uchendu I, Hewitt-Taylor J, Nwakasi C (2021) Knowledge, attitudes, and perceptions about cervical cancer, and the uptake of cervical cancer screening in Nigeria: An integrative review. Sci African 14: e01013.
21. Landis FC, Fai NY, Enow-Orock G, Omer N MN (2022) Perception on cervical cancer and enhancing strategies to screening uptake among women in Bamenda, Cameroon: A qualitative method. J Public Heal Epidemiol 14: 149-159.
22. Burrowes S, Holcombe SJ, Leshargie CT, Hernandez A, Ho A, et al. (2022) Perceptions of cervical cancer care among east Gojjam zone women and their providers: a qualitative study. Reprod Health 2: 1-18.