

Acute Fulminant Pancreatitis in a Patient with Excessive Erythrocytosis

Arturo Zárate Curi^{1,2}, Luis Enrique Núñez Moscoso^{1,2*}, Brenda Caira¹, Karoline Zarate Pareja³, Alejandro Ramos Paredes¹ & Ivan Lira Portugal^{1,2}

¹Emergency Department, Hospital Nacional Carlos Alberto Seguin Escobedo-EsSalud, Arequipa, Peru

²Medicine Faculty, Universidad Nacional de San Agustín, Arequipa, Peru

³Biotechnology Engineering School, Universidad Católica de Santa María, Arequipa, Peru

*Corresponding author: Luis Enrique Núñez Moscoso, Emergency Department, Hospital Nacional Carlos Alberto Seguin Escobedo-EsSalud, Arequipa, Peru.

Submitted: 20 February 2024 Accepted: 26 February 2024 Published: 04 March 2024

doi <https://doi.org/10.63620/MKWJMHC.2024.1008>

Citation: Zárate Curi, A., Núñez Moscoso, L. E., Caira, B., Zarate Pareja, K., Ramos Paredes, A., & Lira Portugal (2024) Acute Fulminant Pancreatitis in a Patient with Excessive Erythrocytosis. *Wor Jour of Medic and Heal Care* 2(1), 01-04.

Abstract

Introduction: Severe acute pancreatitis is associated with a high morbidity and mortality due to the development of pancreatic and extrapancreatic necrosis, and subsequently necrotic tissue infection resulting in multisystem organ failure. Excessive erythrocytosis and hyperviscosity could promote hypoperfusion and ischemia of pancreatic tissue, favoring the systemic inflammatory response (SIRS) and worsening the course of the acute pancreatitis.

Case Presentation: We describe the case of a 71-year-old male patient from Arequipa, who worked in La Riconada, Puno, the place with the highest altitude above sea level (5300 msnm) in Peru. The patient presented to the emergency department with several abdominal pain and dyspnea, the physical examination revealed peritoneal irritation and cyanosis in the extremities, and analysis revealed a possible coagulation disorder. The patient presented with severe acute pancreatitis along with acute respiratory failure requiring an intensive care unit for management. Extreme polycythemia and insufficient management quickly led to necrotizing pancreatitis that led to multisystem failure and death of the patient.

Conclusion: In conclusion, we report a case of severe acute pancreatitis in a patient with extreme polycythemia that rapidly led to necrotizing pancreatitis, multiorgan failure and patient death, which is not commonly reported in the literature in the management of patients diagnosed with severe pancreatitis.

Keywords: Accute Severe Pancreatitis, Polycythemia, Peru

Introduction

Globally, it is estimated that more than 140 million people live in localities above 2500 meters above sea level (masl), of these, 80 million people live in Asia and 35 million in the Andes, in the later the highest population density was found above 3500 meters above sea level [1, 2]. The population at this altitude lives in an environment of hypobaric hypoxia and consequently low partial pressure of inspired oxygen. As a result, he develops chronic alveolar hypoxia, hypoxemia, and polycythemia [3].

According to the International Consensus on Chronic Diseases published in 2005, Chronic Mountain Sickness is characterized by excessive erythrocytosis (women Hb \geq 19 g/dL; men Hb \geq 21 gr dL), severe hypoxemia and in some cases, moderate or severe pulmonary hypertension which can progress to pulmonary heart and heart failure, due to maladaptation to altitude, in habitants at more than 2500 meters above the sea level [4].

Most patients with acute pancreatitis develop a mild course, but about 15-20% progress to severe acute pancreatitis, defined as the presence of organ failure or local complications such as necrosis, abscess or pseudo-cyst which is associated with a mortality rate of 3-5% despite advances in its management [5].

In many cases, inadequate diagnostic approach in patients with a significant epidemiological history may lead to inadequate clinical management and thus probably increase morbidity and mortality and the use of unnecessary medical resources.

The objective of this report is to present a case of severe acute pancreatitis with fatal outcome in a male patient with concomitant excessive erythrocytosis, inhabitant of one of the highest villages in Peru and the world (La Rinconada, Puno at 5300 m.a.s.l.); to subsequently carry out a review of the literature pertinent to the case.

Case Presentation

Male patient, 71 years old. No history of chronic diseases. He worked in an informal mine located in La Rinconada, district of Ananea in the province of Puno, which is 5300 meters above the sea level (masl) and therefore considered the highest town in the world, approximately every 6 months he descended to a lower altitude, like Arequipa. Likewise, he denied undergoing periodic medical evaluations.

He went to the Emergency Department of the Carlos Alberto Seguin Escobedo National Hospital in the city of Arequipa, reporting an apparently 12-hours sick time, whose main symptom was acute abdominal pain after ingestions of fatty foods, located in the epigastrium and radiated to the dorsum, of great intensity (Analog Pain Escala 10/10) oppressive type, accompanied by nausea, vomiting at various times and sweating. He denied having previously self-medication. He denied the intake of alcoholic beverages chronically and prior to the onset of the disease.

We were struck by the poor general condition with respiratory, neurological and hemodynamic compromise with which he was admitted.

Admitted in poor general condition, sporadic, Glasgow Scale 10/15, dry oral mucosa, capillary refill in more than 3 seconds, blood pressure 100/60 mmHg, HR 98/min, SpO₂ 75% with 0.21 and 80% with FiO₂ 90% (mask with reservoir), RR 40/min, with poor ventilatory pattern, tachypneic, drowsy, cold skin and mild cyanosis in fingernails and toenails, the abdomen was distended, painful to superficial and deep palpation, with signs of peritoneal irritation (Blumberg and Rovsing positive) and intestinal ileus.

Regarding laboratory tests, the antigenic and molecular test for COVID-19 were negative, leukocytosis 12,370/mm³, Hemoglobin 27.4 mg/dL, Hematocrit 81.7%, Platelets 120,000/uL, glucose 298 mg/dL, total bilirubin 3.4, direct bilirubin 1.8, prothrombin time 26.7 seconds, INR 2.52, TPT 86.6 segundos, TGO 234 U/L, TGP 148 U/L, amylase 1455 U/L, lipase 1420 U/L, CRP 19.7 mg/dL, Urea 54.6 mg/dL, creatinine 2.07 mg/dL, Ph 7.25, PCO₂ 38, PO₂ 70 mmHg, HCO₃ 17 mmHg, lactate 6.7 mmol/L, Na 133 mmol/L, K 4.3 mmol/L, Ca 1.11 mmol/L, Cl 112 mmol/L, D-dimer 1.35 ug/mL, troponin T 0.003 ng/mL, CPK 68 U/L, proBNP 3263 pg/mL Apache II in 22 points with a calculated mortality of 54.9%, BISAP of 04 points and calculated mortality higher than 15%.

The chest tomography (TEM) showed condensation-type hyperdensities in both bases and pleural effusion and the abdominal TEM showed enlarged pancreas with acute inflammatory signs and small collections and gallbladder with absence of lithiasis, concluding: Acute necro-hemorrhagic pancreatitis.

Echo cardiography: severely depressed left ventricular (LV) systolic function, LVEF 20%, basal and medial LV akinesia.

He was intubated and connected to invasive mechanical ventilation, received support with active vessel amines and others for a critically ill patient.

A detail observed at the time of performing the hemogluco-test was that the blood coagulated quickly, making it difficult to register it.

16 hours after being admitted to the Emergency Department, he was transferred to the ICU. At no time was hemodynamic stabilization achieved, but rather greater Multiorgan Dysfunction (MOD), requiring higher doses of amines (Norepinephrine, Vasopressin, Dobutamine) with Acute Renal Failure, Acute Respiratory Failure with Acute Respiratory, Hepatic, Gastrointestinal and Cardiovascular Distress with acute stress cardiomyopathy with Takotsubo syndrome and therefore unfavorable prognosis in the short term, dying 30 hours after admission.

Discussion

The patient came from La Rinconada, located in the department of Puno, in the south-eastern region of Peru, considered to be the highest town in the world with more than 17,000 inhabitants, located at an altitude of up to 5300 meters above sea level who developed severe and fulminant acute pancreatitis, it was not possible to demonstrate the two main causes of it such as gallstones and alcohol, but rather we were very surprised to find one of the highest values of Hemoglobin (Hb) and Hematocrit (Hto) [6, 7].

In relation to the abnormal elevation of Hb and Hto, there is terminology that needs to be clarified. High-altitude erythrocytosis differs from terms such as secondary polycythemia and polycythemia vera, which would be related to chronic diseases such as COPD, pulmonary fibrosis, etc. or an onco-hematological disease in which there is an alteration not only in the concentration of haemoglobin but also of the three hematopoietic series, respectively [8].

In relation to the term acute fulminant pancreatitis, the following was found in the literature: the criteria for defining early severe AP (ESAP) was severe organ failure (OF) within 07 days after pancreatitis. Patients with ESAP were subdivided into fulminant and subfulminant AP according to the time of OF, i.e., <72 h and between 4 and 7 days of pancreatitis, respectively, it is also described that the greater the organ failure, the higher the mortality [9].

The patient developed severe acute pancreatitis according to the Atlanta classification, and the Apache II and BISAP scores, the systemic inflammatory response was aggressive and progressive towards multiple organ failure, requiring ventilatory support and hemodynamic monitoring [10-13].

In AP, the reduction in intravascular volume can be detected as an increase in the serum hematocrit level, which can lead to decreased perfusion of the microcirculation of the pancreas and result in pancreatic necrosis. In a study conducted by Banks on "Hemoconcentration as an Early Predictor of Organ Failure and Pancreatic Necrosis" they concluded that a hematocrit greater than or equal to 44 at admission or during the first 24 hours had an increased risk of pancreatic necrosis and that a decrease in hematocrit during the first 24 hours had a lower risk of pancreatic necrosis however, most studies show that this is not a reliable predictor of severity [14-17].

Excessive high altitude erythrocytosis would be an important factor in the development of severe acute pancreatitis pathologically speaking. Blood concentration and blood hyperviscosity is a risk factor for the severity of acute pancreatitis as described by multiple studies such as severity prognostic factors in patients with acute pancreatitis in hospital nacional sergio e. vernaes [18]. In its conclusions, it was found that patients with a hematocrit >40% were 4.17 times more likely to develop moderately severe acute pancreatitis and severe acute pancreatitis, as opposed to those who obtained a hematocrit <40% (OR: 4,17; IC 95%: 2,07-8,39; value p: <0,001)

Hemoconcentration would be due to multiple causes, such as the implicit state of volume depletion due to lack of intake, vomiting, abnormal distribution of water in the third space, etc. Hematocrit values above 40% have been described related to severity and higher mortality from this disease, this in turn would condition less perfusion at the level of the microcirculation of the pancreatic tissue, with a risk of ischemia and necrosis, clinically expressed in severe presentation of acute pancreatitis and increased risk of mortality [19].

The presence of SIRS and its duration over time would be related to a greater severity of acute pancreatitis.

In the study by Hai-Hong Zhu et al, called The relationship between systemic inflammatory response syndrome and severity of acute pancreatitis combined with plateau erythrocythemia, they found that there was a significant difference between the erythrocythemia group and the non-erythrocythemia group, not only in the incidence of patients who patients who developed SIRS, but also in two elements of patients who did or did not meet the diagnostic criteria for SIRS ($P < 0.05$). The more severe the acute pancreatitis combined with erythrocythemia, the longer the duration of SIRS [20-22].

More recently (2022), the same researcher in his publication, Association of High Altitude Polycythemia with an Increased Risk of Systemic Inflammatory, Response Syndrome in Acute Pancreatitis describes a retrospective study evaluating the relationship between acute pancreatitis and high-altitude polycythemia in 100 patients admitted to the People's Hospital of Qinghai Province during 2006-2016. Patients were divided into two groups: one with acute pancreatitis and high-altitude polycythemia (PA+HAPC) and one with acute pancreatitis without high altitude polycythemia (AP). The results showed that high-altitude polycythemia was associated with an increased risk of SIRS in patients with acute pancreatitis. In addition, the greater the severity of acute pancreatitis combined with high-altitude polycythemia, the longer the duration of SIRS. The development of SIRS and the severity of AP+HAPC are closely related. Patients with HAPC are more vulnerable to the effects of AP with concomitant SIRS [23].

Multiorgan dysfunction in severe pancreatitis has been reported to be greater in patients with hypoxia due to high-altitude erythrocytosis. In the studio The changes and significance of multiple organ functions in acute pancreatitis patients under hypoxic condition on plateau, Hai-hong ZhuL, describes that alanine aminotransferase (ALT) and creatinine (Cr) were significantly higher

in patients with AP who complicated plateau erythrocythemia compared with uncomplicated AP patients with plateau erythrocythemia. Arterial partial pressure of oxygen (PaO₂) and arterial oxygen saturation (SaO₂) were significantly lower in AP with plateau erythrocythemia complication than AP without complication of plateau erythrocythemia. There was no difference in aspartate aminotransferase (AST), blood urea nitrogen (BUN) and partial blood pressure carbon dioxide (PaCO₂); however, their levels were higher in cases of plateau erythrocythemia than in those without plateau erythrocythemia [24].

Conclusions

In conclusion, we report the case of a 71-year-old male patient with excessive erythrocytosis who developed severe acute pancreatitis. Erythrocytosis possibly accelerated the picture rapidly leading to necrotizing pancreatitis, multi-organ failure and death of the patient, this is not commonly reported in the literature in the management of patients diagnosed with severe pancreatitis. Clinical research in patients with chronic conditions such as erythrocytosis is essential to establish the clinical approach and management of these patients in urgent or emergency settings such as acute pancreatitis or others. In addition to recognizing that timely diagnosis and treatment is essential to avoid severe complications that quickly progress.

Contribution to Authorship

All authors writing of the manuscript, and all participated in the review and approval of the final manuscript.

Funding Statement

Self-funded

Disclosure

The authors declare no competing conflicts of interests.

References

1. Moore, L. G., Niermeyer, S., & Zamudio, S. (1998). Human adaptation to high altitude: Regional and lifecycle perspectives. *American Journal of Physical Anthropology*, 27, 25-64.
2. Eichstaedt, C. A., Antao, T., Pagani, L., Cardona, A., Kivisild, T., & Mormina, M. (2014). The Andean adaptive toolkit to counteract high altitude maladaptation: Genome-wide and phenotypic analysis of the Collas. *PLoS ONE*, 9, e93314.
3. Peñaloza, D., & Arias Stella, J. (2011). Heart and pulmonary circulation at high altitudes: Normal natives and chronic mountain sickness. *Peruvian Journal of Cardiology*, 37, 38-56.
4. León-Velarde, F., Maggiorini, M., Reeves, J. T., Aldashev, A., Asmus, I., Bernardi, L., ... & Zubieta-Calleja, G. (2005). Consensus statement on chronic and subacute high-altitude diseases. *High Altitude Medicine & Biology*, 6, 147-157.
5. McKay, C. J., Evans, S., Sinclair, M., Carter, C. R., & Inrie, C. W. (1999). High early mortality rate from acute pancreatitis in Scotland, 1984-1995. *British Journal of Surgery*, 87, 1302-1305.
6. West, J. B. (2004). Highest permanent human habitation. *High Altitude Medicine & Biology*, 3.
7. Correo. (2016, October 25). La ciudad más alta del mundo y sombría está ubicada en Puno [The highest and most

- gloomy city in the world is located in Puno]. <https://diarioocorreio.pe/edicion/puno/la-ciudad-mas-alta-del-mundo-y-sombrio-esta-ubicado-en-puno-video-696750/>
8. Uscamayta, N. (2007). Pathological high erythrocytosis. Scientific. Reviewed on the Internet, 5, 50-56.
 9. Sharma, M., Banerjee, D., & Garg, P. K. (2007). Characterization of newer subgroups of fulminant and subfulminant pancreatitis associated with a high early mortality. American Journal of Gastroenterology, 102, 2688-2695.
 10. Petrov, M. S., & Windsor, J. A. (2010). Classification of the severity of acute pancreatitis: How many categories make sense? American Journal of Gastroenterology, 105, 74-76.
 11. Banks, P. A., Bollen, T. L., Dervenis, C., Gooszen, H. G., Johnson, C. D., Sarr, M. G., ... & Vege, S. S. (2013). Classification of acute pancreatitis—2012: Revision of the Atlanta classification and definitions by international consensus. Gut, 62, 102-111.
 12. Knaus, W. A., Draper, E. A., Wagner, D. P., & Zimmerman, J. E. (1985). APACHE II: A severity of disease classification system. Critical Care Medicine, 13, 818-829.
 13. Wu, B. U., Johannes, R. S., Sun, X., Tabak, Y., Conwell, D. L., & Banks, P. A. (2008). The early prediction of mortality in acute pancreatitis: A large population-based study. Gut, 57, 1698-1703.
 14. Vege, S. S., Gardner, T. B., Chari, S. T., Munukuti, P., Pearson, R. K., Clain, J. E., ... & Sarr, M. G. (2009). Low mortality and high morbidity in severe acute pancreatitis without organ failure: A case for revising the Atlanta classification to include “moderately severe acute pancreatitis.” American Journal of Gastroenterology, 104, 710-715.
 15. Wu, B. U., Johannes, R. S., Conwell, D. L., & Banks, P. A. (2009). Early hemoconcentration predicts increased mortality only among transferred patients with acute pancreatitis. Pancreatology, 9, 639-643.
 16. Brown, A., Orav, J., & Banks, P. A. (2000). Hemoconcentration is an early marker for organ failure and necrotizing pancreatitis. Pancreas, 20, 367-372.
 17. Chatzicostas, C., Roussomoustakaki, M., Vlachonikolis, I. G., Notas, G., Mouzas, I., Samonakis, D., & Kouroumalis, E. A. (2002). Comparison of Ranson, APACHE II and APACHE III scoring systems in acute pancreatitis. Pancreas, 25, 331-335.
 18. Jamanca-Milian, H., & Cano-Cardenas, L. (2020). Prognostic factors of severity in patients with acute pancreatitis in the gastroenterology service of the Sergio E. Bernales National Hospital. Revista de la Facultad de Medicina Humana, 20, 14-19.
 19. Sanchez, C. (2015). Value of hemoconcentration as an early predictor of severity in acute pancreatitis, in patients admitted to Hospital México during the period from January 2003 to December 2012 [Master's thesis, University of Costa Rica]. Sistema de Estudios de Posgrado.
 20. Zhu, H. H., Wu, X. M., Guo, Y. M., Yang, J. Y., Yexie, Z. H., Ye, C. J., & Bai, Y. L. (2010). The relationship between systemic inflammatory response syndrome and severity of acute pancreatitis combined with plateau erythrocythemia. Revista China de Cirugía, 48, 1137-1140.
 21. Cubas, S., Varela, M., Noria, A., Ibarra, S., Martínez, J. P., González, F., & Canessa, C. (2017). SIRS as a predictor of severity in acute pancreatitis. Revista Médica del Uruguay, 33, 174-179.
 22. Du, W., Wang, H., Zhang, S. W., & Wang, B. E. (2005). Investigation on the relation between systemic inflammatory response syndrome and severity of acute pancreatitis. Zhongguo Wei Zhong Bing Ji Jiu Yi Xue, 17, 279-281.
 23. Zhu, H.-H., Guo, Y.-M., Yang, J.-Y., Yexie, Z.-H., Yan-Lin. (2022). Association of high-altitude polycythemia with an increased risk of systemic inflammatory response syndrome in acute pancreatitis. SL Gastroenterology, 4, 126.
 24. Zhu, H. H., Wu, X. M., & Ye, C. J. (2010). The changes and significance of multiple organ functions in acute pancreatitis patients under hypoxic condition on plateau. Zhongguo Wei Zhong Bing Ji Jiu Yi Xue, 22, 210-213.