

Ileoileal Knotting a Rare Cause of Double Loop Small Bowel Obstruction, Diagnostic and Intraoperative Challenge at Ethiopian Leku District Hospital, Sidama Region, April 2023, Case Report

Asfaw Kibret

leku hospital, sidama, Ethiopia

*Corresponding author: Asfaw Kibret, leku hospital, sidama, Ethiopia

Submitted: 20 June 2023

Accepted: 27 June 2023

Published: 30 June 2023

doi <https://doi.org/10.63620/MKJCSCR.2023.1005>

Citation: Kibret, A. (2023). Ileoileal Knotting a Rare Cause of Double Loop Small Bowel Obstruction, Diagnostic and Intraoperative Challenge at Ethiopian Leku District Hospital, Sidama Region, April 2023, Case Report. *J Clin surg Care Res*, 2(2), 01-03.

Abstract

Background: Intestinal obstruction is common cause for abdominal surgeries. Among the causes for intestinal obstruction ileal knotting is the unusual and rare cause with very few case reports in the globe and in Ethiopia. Almost all cases were intraoperative surprises. We report a case of strangulated closed small bowel loop obstruction caused by ileoileal knotting at distal ileum near to ileocecal junction, its diagnostic and management challenges and the role of the “**Integrated Emergency surgical officers (IESO)**” for early emergency surgical interventions in district primary hospitals in Ethiopia where there is scarce general specialist surgeon.

Case Presentation: Here we discuss a 78yrs old male Ethiopian patient presented to our district hospital with sign symptoms of strangulated small bowel obstruction and exploratory laparotomy was done with intraoperative surprise of ileoileal knot with gangrenous closed loop obstruction at ileocecal junction. The gangrenous segment was resected and ileo-ascending colon anastomosis was done. The post-operative stay was uneventful and patient discharged on his 7th post operation date.

Conclusion: Here we present this case to highlight difficulty of preoperative diagnosis and the need of high index of suspicion as well as the management challenges of resection anastomosis when the knot is near to ileocecal junction and the undeniable role of emergency surgical officers for early interventions of such emergency cases who need early interventions.

Keywords: Closed Loop Obstruction, Gangrenous Bowel, Emergency Surgical Officers, Ileoileal-Knotting

Introduction

Intestinal obstruction is common cause of acute abdomen which needs emergency surgery among which Small bowel obstruction is the commonest. There are different causes of SBO which includes primary volvulus, hernias, adhesions, intussusceptions and intestinal knotting. Intestinal knotting are the rare causes for bowel obstructions. Ileosigmoid knotting is the commonest and ileoileal knotting is the unusual and only reported in few cases [1, 2].

District hospitals in Ethiopia are primary level hospitals and usually the surgical unit is lead by Integrated Emergency surgical officers (IESO) who is dully known in Ethiopia and the IESO is a postgraduate professional trained for 3yrs and authorized to do emergency obstetrics, gynecologic and emergency general surgeries. It was task shifting professional license allowing to do

emergency surgeries in district Ethiopian hospitals where there are scarce physician specialists [3].

According to reports showing the clinical performance of IESOs there are numerous emergency surgeries has been doing in each day in district hospitals in Ethiopia. Emergency laparotomies are the common procedures done and SBO is the commonest one during emergency bowel surgeries. Adomen is a “Pandora’s box” till it is opened which later comes with intraoperative surprises specially in developing country like Ethiopia where there is no essential diagnostic imaging in district hospitals. In this article we present a 78yrs old male patient who presented with signs and symptoms of small bowel obstruction and diagnosed intraoperatively to have double loop obstruction secondary to ileoileal knotting. We will present the diagnostic and intraoperative challenges of the case [3].

Case Presentation

This is 78yrs old male farmer presented to Leku district hospital emergency department with history of abdominal pain of 28hrs duration. The pain was central and intermittent colicky type initially. He had also associated history of repeated bilious type vomiting. He had history of failure to pass feces and flatus since the last one day before his presentation to hospital, later he develops abdominal distention and the pain becomes persistent and generalized. He had no any history of previous abdominal surgery, no history of abdominal trauma. He has no history of diagnosed hypertensive, cardiac and diabetes mellitus diseases. Upon presentation on his physical examinations he was acute sick looking with vital signs of BP=90/50mmgh, PR=129bpm, RR=23br/min, T=36.9, Pso2=96% at atmospheric air. He had dry tongue and dry buccal mucosa, with little eye sickening. On abdominal examination he had distended abdomen, hyper tympanic on percussion, there was direct tenderness over his abdomen upon palpation, on Digital rectal examination the rectum was empty. CBC and blood group, were determined with the result of WBC=12.4, HCT=28.3, PLT=258, BG=BgRh=O+, Erect abdominal X-ray was sent and showed dilated small bowel loops with multiple air fluid levels (Fig.1)

Based on the above findings with presumed diagnosis of generalized peritonitis secondary to Strangulated SBO secondary to small bowel volvulus, exploratory laparotomy was decided and the patient resuscitated by 3bags of normal saline. NG tube

was inserted, broad spectrum parenteral antibiotics was started. Lately with preoperative vital signs of BP=110/60, PR=102bpm, RR=23br/min, Pso2=96%, UOP =200ml/1hr, anesthesia team was consulted for preoperative anesthesia evaluation and evaluated. then taken to OR. Under general anesthesia the patient abdomen has been opened via the midline incision. Upon entry to the general peritoneal cavity there was offensive dark hemorrhagic fluid estimated to be 800ml. There were dilated small bowel loops and gangrenous segment was at the distal ileum near to ileocecal junction (ICJ) with ileoileal knotting causing closed loop strangulated obstruction. (Fig.2). Then the knotted bowel was tried to be unraveled but it was difficult. Later it was found that ligating and resecting the gangrenous bowel mesenteric vessels was access able for proximal resection of the gangrenous part of the bowel loops and controlled decompression was done. Then after decompression the bowel ends tied and this enables to unravel and exposed the distal end of the gangrenous segment which also has been resected. The distal aspect of the bowel was near the ICJ which made end to end ileoileal anastomosis difficult. Then ileo ascending colon side to side anastomosis was done in two layers using 2-0 vicryl. Intraoperatively the patient was hypotensive and managed. Finally, the general peritoneal cavity was lavaged by warm saline and abdomen closed layer by layer. Patient transferred to recovery stable, and the post-operative course was uneventful, sips was started on 3rd post operative date and soft regular diet on the next day, discharged stable on 7th post operation date.




Figure 1: Erect abdominal x-ray with dilated small bowel loops and multiple air-fluid levels



Figure 2: The intra operative finding showing the knot, the non-viable segment and dilated small bowel loops

Discussion

Intestinal obstruction is common cause for emergency surgery and small bowel obstruction is the leading cause for it. There are many causes for SBO. The common causes for SBO in the world and in our country are small bowel volvulus, adhesions, hernia and intussusceptions. Bowel knots are the unusual causes for bowel obstruction among others ileo-ileal knotting are the rarest reported causes for SBO. In my knowledge and review only 14 such cases were reported worldwide and only 3 cases in Ethiopia all of which were managed in specialized hospitals and managed by specialist physicians [1-3]. But in our case, it was managed in one of our district hospitals in Ethiopia and the leading surgeon was IESO who is a holder of MSc in integrated emergency surgery, dully known in Ethiopia and licensed to do obstetrics, gynecology and general surgical emergencies. For last 10yrs IESOs in Ethiopia were performing tremendous emergency surgical activities in primary health institutions in the country where still the physician specialists are scarce [3].

Surgical scholars well described abdomen as “Pandora box” which is most time difficult to speculate what is in unless it is opened specially in developing setups where there is no sophisticated diagnostic imaging’s like standard ultrasounds and CT scan.

Most of the time intestinal knotting is diagnosed intraoperatively. Preoperative diagnosis is very difficult and almost impossible. Ileioleal knotting present like most other small bowel obstruction cases and has no typical or classic sign and symptoms to it except rapid progression to bowel strangulation due to its closed loop obstruction natures. Early operative intervention is mandatory after vigorous resuscitation, nasogastric decompression and broad-spectrum parental antibiotics. Emergency laparotomy should be performed through the midline incision and meticulous exploration of the bowel should be performed. Specific intervention depends on the site of the obstructed knot, the viability of the bowel loops, the length of distal bowel segment away from Ileocecal junction. The choice of operative technique includes careful unraveling the knot if both loops are viable, performing enblock resection if it is found gangrenous and easily accessible the distal and proximal segments. In articles controlled decompression and untie the knot is also suggested. But this may have risk of spillage peritoneal contamination. However, in our case the knot was at distal ileum near to ICJ difficult to enblock resection at distal segment initially the mesentery is ligated and resected and proximal segment crushed resected after control decompression, the decompressed bowel end is tied by sure and easily the knot released and unraveled, the distal part accessed near to ICJ, then ileo ascending anastomosis was done [4-8].

Conclusion

Ileioleal bowel knotting is a rare cause for SBO but it should be among the differentials, it needs high index of suspicion. Its specific operative management depends on the bowel segment accessibility, viability and site of obstruction and mesentery resection and resection of the proximal segment to decompress is possible if the distal is not accessible and if it is near to ICJ. Emergency surgical officers also have undeniable role in the surgical intervention of cases which need emergency operations

Declarations

Ethical Approval

certify that the study was performed in accordance with the ethical standards as laid down in the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards. Leku hospital ethics committee at leku hospital reviewed and approved for submission of the manuscript.

Consent to Participate

Informed consent was obtained from all participants and from a parent and/or legal guardian. Written

Consent for Publication

Written informed consent was obtained from the patient for publication of this study and accompanying images.

Competing Interest

No competing interest.

Authors Contribution

Me as corresponding author and ESO did the operation, prepare the manuscript.

Funding

No funding

Availability of Data and Materials

The patient data and materials are available in the institution (the hospital) data room.

References

1. Abebe, E., Asmare, B., & Addise, A. (2015). Ileo-ileal knotting as an uncommon cause of acute intestinal obstruction. *Journal of Surgical Case Reports*, 2015(8), 1–2.
2. Fekadu, G., Tolera, A., Bayissa, B. B., Merga, B. T., & Edessa, D., et al. (2022). Epidemiology and causes of intestinal obstruction in Ethiopia: A systematic review. *SAGE Open Medicine*, 10, 20503121221083207.
3. Gobeze, A. A., Kebede, Z., Berhan, Y., & Ghosh, B. (2016). Clinical performance of emergency surgical officers in Southern Ethiopia. *Ethiopian Journal of Health Sciences*, 26(5), 463–470. <https://doi.org/10.4314/ejhs.v26i5.6>
4. Kanamori, K., Koyanagi, K., Hara, H., Nakamura, K., Yamamoto, M., et al. (2021). Small bowel obstruction caused by a true ileo-ileal knot: A rare case successfully treated by prior ligation of mesenteric vessels. *Surgical Case Reports*, 7, 195.
5. Beg, M. Y., Bains, L., Lal, P., Maranna, H., & Kumar, N. P. (2020). Small bowel knots. *Annals of the Royal College of Surgeons of England*, 102(7), 571–576.
6. Mohammed, Y., & Tesfaye, K. (2021). Ileioleal knotting: A rare cause of intestinal obstruction: A case report. *Journal of Medical Case Reports*, 15, 397. <https://doi.org/10.1186/s13256-021-03083-y>
7. Sohaila, M., Alyson, T., Sim, S. K., & Nik Azim, N. A. (2020). Intestinal knotting: A case report and brief literature review. *Medical Journal of Malaysia*, 75(6), 606–608.
8. Taniguchi, K., Iida, R., Watanabe, T., Nitta, M., Tomioka, M., et al. (2017). Ileo-ileal knot: A rare case of acute strangulated intestinal obstruction. *Nagoya Journal of Medical Science*, 79(1), 109–113.