

Screening of Visual Pathway in Patients Suffering from Guillain Barre by Visual Evoked Potential

Fatemeh Sarzaeim¹, Fatemeh Aflaki², Seyed Mohammad Masoud Shushtarian^{2*}, and Ahmad Shojaei³

¹Department of Neurology, Faculty of Medicine, Tehran Medical Sciences, Islamic Azad University, Tehran, Iran.

²Department of Biophysics and Biochemistry, Faculty of Advance Science and Technology, Tehran Medical Sciences, Islamic Azad University, Tehran, Iran.

³Basir Eye Health Research Center, Iran University of Medical Sciences, Tehran, Iran.

***Corresponding author:** Seyed Mohammad Masoud Shushtarian, Department of Biophysics and Biochemistry, Faculty of Advance Science and Technology, Tehran Medical Sciences, Islamic Azad University, Tehran, Iran.

Submitted: 08 April 2024 Accepted: 15 April 2024 Published: 22 April 2024

doi <https://doi.org/10.63620/MKSSJOEC.2024.1009>

Citation: Sarzaeim, F., Aflaki, F., Shushtarian, S. M. M., & Shojaei, A. (2024). Screening of Visual Pathway in Patients Suffering from Guillain Barre by Visual Evoked Potential. *Sci Set J of Ophthalmology & Eye Care*, 3(2), 01-04.

Abstract

Aim: Guillain-Barre syndrome (GBS) is a rare condition in which a person's immune system attacks the peripheral nerve. The visual system can also be affected in GBS patients. The aim of present work is to look for visual pathway disturbances in these patients using visual evoked potential (VEP).

Material and Methods: 10 male patients (20 eyes) diagnosed as GBS in age range of 25 to 56 years were selected for the purpose of present study. Visual evoked potential (VEP) using pattern type of stimulation was tested in total patient group using Mangoni machine. The results obtained was compared with 10 normal sex and age matched population following VEP test.

Results: There was not statistically significant differences as far as demographical finding was concerned between case and control groups.

There was a significant difference was observed in latency and amplitude of VEP, P100 Peak between case and control groups ($P < 0.001$ for both parameters).

Conclusion: Guillain barre syndrome can affects the visual pathway of visual system which can be measured by VEP.

Keywords: Guillain-Bare Syndrome, Visual Pathway, Visual Evoked Potential

Introduction

Guillain-Barre syndrome (GBS) is a rare autoimmune-mediated peripheral neuropathy with an acute onset. GBS is associated with posterior cranial nerve injury including the facial, oculomotor, glossopharyngeal and vagus nerve. Many parts of the body including visual system can be affected in these patients. Electrophysiological techniques of the vision are used to assess different pathological conditions of visual system, mainly the retina and visual pathway. Electroretinography (ERG), electrooculography (EOG) and visual evoked potential (VEP) are commonly employed electrophysiological techniques used in this field.

Abdolalizadeh S et al (2022) conducted A study to investigate the potential effects of antiseizure medication on patients using

ERG. The study included twenty participants consisting of ten males and ten females, ranging in age from 15 to 30 years. The findings revealed retinal changes in these patients which were diagnosed by measuring the amplitude of ERG, specifically b-wave peak [1]. The same research team also examined the retinal pigment epithelium (RPE) of patients undergoing treatment with anti-epileptic medication using EOG. They utilized the same group of patients and observed pathological changes in the RPE, which were identified by assessing the Arden Index (AI) of EOG test [2]. Shushtarian S M et al (2017) designed a study to investigate the potential effects of vibration on the visual pathway using VEP. They selected 50 workers from a textile factory segment where machinery creating high level of vibration. The study concluded that occupational vibration had adverse effects

on the visual pathway, leading to increased latency of VEP, specifically the P100 Peak [3]. Numerous references have been published on this topic [4-47].

In present study we utilized VEP to screen the visual pathway of the patients who had suffered from GBS.

Material and Methods

In a case control study, we selected 10 male patients (20 eyes) with GBS as the case group. The patients in the case group were within the age range of 30-65 years. VEP test was examined in total case group. For stimulating the eyes of patient's pattern reversal checkboard was used. Latency (msec) and amplitude (μ V) of VEP, P100 Peak was measured. As a comparison group, we

selected 10 healthy subjects (visual pathway) of same age and sex of case group, as a control group. For recording VEP in both the group we used to connect the total population to the recording machine, i.e., active, reference and earth were attached to occipital region, vertex and forehead respectively. The setup was designed to ensure accurate and reliable measurements during VEP procedure.

Results

Table 1, Comparison of the mean age in the case and control groups. There were no statistically significant differences between the two groups in terms of age ($P=0.492$). and sex (all participants were male).

Table 1: Demographical finding in case and control groups.

Variable	Number of participants	groups (Mean \pm SD)		P value*
			Case	
Age	10 (20 eyes)	45.61 \pm 6.8	45.72 \pm 7.12	0.492

* Based on Mann-Whitney Test

Table 2 presents a comparison of latency and amplitude measurements for the VEP P100 peak between the case and control groups. It was observed that the case group had a statistically higher latency ($P < 0.001$) and lower amplitude for the VEP P100 Peak ($P < 0.001$) compared to control group.

Table 2: Comparison of mean latency and amplitude measurements of VEP, P100 peak in the case and control groups.

Variable	Number of participants	groups (Mean \pm SD)		P value*
		Control	Case	
Latency (msec)	10	97.46 \pm 2.8	108.92 \pm 3.84	0.000
Amplitude (μ V)	10	6.38 \pm 1.6	2.96 \pm 1.12	0.000

* Based on Mann-Whitney Test

Discussion

Guillain-Barre syndrome (GBS) can damage the visual pathway of the patients which can be measured by latency and amplitude of VEP, P100 Peak. It is a fact that VEP is a suitable technique to screen the visual pathway [48-50].

The findings of present work may be supported by following research work.

Güngör L et al (2011) worked on 32 patients with diagnosis of GBS. They recorded VEP in total patient group. In their work only five cases (16%) had abnormal VEPs, i.e., mainly increase in latency of VEP, P100 Peak. Zgorzalewicz H et al (2004) performed an extensive research work on VEP in children and adolescents with GBS [51]. They found a few patients had abnormal VEP with prolongation of latency of VEP, P100 Peak with changes of amplitude of same peak [52].

These two works support the findings of present work to some extent however in two mentioned work a few GBS had abnormal VEP where as in our work maximum number of patients had abnormal VEP's. The reason for this difference lies in the fact that the other research workers used random patients where as we consider the GBS patients who referred for VEP test with probability of visual pathway deficits [53].

Conclusion

Guillain- Barre syndrome can affect the visual pathway of human visual system and can be measured by latency and amplitude of visual evoked potential P100 Peak.

Conflict of Interest

The authors have no conflict of interest with the subject matter of this manuscript.

References

1. Abdolalizadeh, S., Ghasemi, M., Mohammadzadeh, P., Shushtarian, S. M. M., & Shojaei, A. (2022). Retinal Screening of Patients Treated with Antiepileptic Medications Using Electroretinography. *Journal of Ophthalmology and Research*, 5, 165-167.
2. Abdolalizadeh, S., Karami, S., Tava koolpoor Saleh, N., Shushtarian, S. M. M., Pour Mazar, R., et al. (2023). Retinal Pigment Epithelium Screening of Patients Treated with Anti-Epileptic Medications using Electrooculography. *Journal of Ophthalmology and Research*, 6, 8-11.
3. Shushtarian, S. M. M., Shahriyari Kalantari, A., Tajik, F. Â., & Adhami-Moghadam, F. (2017). Effect of occupational vibration on visual pathway measured by visual evoked potentials. *Journal of Ophthalmic and Optometric Sciences*, 1, 7-11.
4. Keramti, S., Ojani, F., Shushtarian, S. M. M., Shojaei, A.,

- & Mohammad-Rabei, H. (2021). Early Diagnosis of Pathological Changes in Visual System of Prolactinoma Patients Using Visual Evoked Potential. *Journal of Ophthalmology and Research*, 4, 289-293.
5. Ojani, F., Shushtarian, S. M. M., Shojaei, A., & Naghib, J. (2021). Visual Evoked Potential Findings of Bardet-Biedl Syndrome. *Journal of Ophthalmology and Research*, 4, 254-257.
 6. Sarzaeim, F., Hashemzehi, M., Shushtarian, S. M. M., Shojaei, A., & Naghib, J. (2022). Flash Visual Evoked Potential as a Suitable Technique to Evaluate the Extent of Injury to Visual Pathway Following Head Trauma. *Journal of Ophthalmology and Research*, 5, 20-23.
 7. Allahdady, F., Amiri, M. A., Shushtarian, S. M. M., Tabatabaee, S. M., Sahrae, F., et al. (2016). Comparison of visual evoked potential and electro-oculogram tests in early detection of hydroxychloroquine retinal toxicity. *Journal of Ophthalmic and Optometric Sciences*, 1, 19-26.
 8. Shushtarian, S. M., Mirdehghan, M. S., & Valiollahi, P. (2008). Retinal damages in turner workers of a factory exposed to intraocular foreign bodies. *Indian Journal of Occupational and Environmental Medicine*, 12, 136-138.
 9. Shushtarian, S. M. M., Mohammad-Rabei, H., & Tahmasebi Barjoui Raki, S. (2018). Effect of Occupational Vibration on Human Retina Measured by Electroretinography. *Journal of Ophthalmic and Optometric Sciences*, 2, 14-17.
 10. Naser, M., Shushtarian, S. M. M., Shojaei, A., & Adlami-Moghadam, F. (2017). Visual Disturbance in a Patient with Amiodarone Treatment Following Refractive Surgery. *Journal of Ophthalmic and Optometric Sciences*, 1, 39-42.
 11. Shushtarian, S. M., Adhami-Moghadam, F., Naser, M., Khazaii, R., Sobhani, S., Shabani, S., et al. (2017). Electroretinographic Changes in Multiple Sclerosis Patients with Abnormal Visual Evoked Potentials. *Journal of Ophthalmic and Optometric Sciences*, 1, 34-38.
 12. Shushtarian, S. M. M., Adhami-Moghadam, F., Naser, M., & Shojaei, A. (2017). Severe Headache Initiated by Flash Stimulation during Visual Evoked Potential Recording in a Patient with Monocular Optic Neuritis and History of Migraine Headache. *Journal of Ophthalmic and Optometric Sciences*, 1.
 13. Keramti, S., Javanshir, S., Tajik, F., Shushtarian, S. M. M., Shojaei, A., et al. (2021). Retinal Screening of Prolactinoma Patients using Flash Electroretinography. *Journal of Ophthalmology and Research*, 4, 321-326.
 14. Hajibeygi, R., Shushtarian, S. M. M., & Abolghasemi, S. (n.d.). Visual Evoked Potential Findings of Sjogren's Syndrome. *Journal of Ophthalmic and Optometric Sciences*, 4, 13-17.
 15. Shushtarian, S. M. M., Adhami-Moghadam, F., Adhami-Moghadam, P., & Abdolhoseinpour, H. (2018). Electrophysiological Eye Examination Changes in a Patient with Sjogren's Syndrome. *Journal of Ophthalmic and Optometric Sciences*, 2, 40-43.
 16. Shushtarian, S. M. M., Tajik, F., & Abdolhoseinpour, H. (2018). Measurement of Visual Page No: 03 Evoked Potentials in Patients with Spastic Cerebral Palsy. *J. Ophthalmic Optom. Sci*, 2, 10-13.
 17. Tajik, F., & Shushtarian, S. M. M. (2018). Electrooculographic and Electroretinographic Changes among Patients Undergoing Treatment with Amiodarone. *Journal of Ophthalmic and Optometric Sciences*, 2, 7-11.
 18. Naser, M., & Shushtarian, S. M. (2014). Study the effect of depakine on retina of epileptic patients using electroretinogram. *International journal of scientific research*, 3, 392-393.
 19. Sarzaeim, F., Ojani, F., Hojati, T. S., Shojaei, A., & Shushtarian, S. M. M. (2022). Effect of Hand-Arm Vibration on Retina of Road Drilling Machine Laborers Measured by Electroretinography. *Journal of Ophthalmology and Research*, 5, 81-85.
 20. Shushtarian, S. M. M., Naghib, S. J., Adhami-Moghadam, F., & Shojaei, A. (2020). Diplopia and Blurry Vision Following Refractive Eye Surgery: A Comorbidity Case Report. *Journal of Ophthalmic and Optometric Sciences*, 4, 40-42.
 21. Shushtarian, S. M. M. (2020). Suitable Stimulation Technique to Record Visual Evoked Potential in Migraine Patients. *Journal of Ophthalmic and Optometric Sciences*, 4, 41-45.
 22. Shushtarian, S. M. M., Pour Mazar, R., & Fadaeifard, S. (2021). Visual Evoked Potential Recording in a Fatigued and Drowsy Patient under Anti-Seizure Medicine Treatment. *Journal of Ophthalmic and Optometric Sciences*, 5, 67-70.
 23. Shushtarian, S. M. M., & Vahid Dastjerdi, M. (2020). Total Blindness Following Anaphylactic Shock due to Co-Amoxiclav Treatment. *Journal of Ophthalmic and Optometric Sciences*, 4, 39-41.
 24. Shushtarian, S. M. M., Haji Naghitehrani, K., & Aflaki, F. (2020). Diplopia and Flashes of Light Sensation in a Patient with Fragrance Allergy. *Journal of Ophthalmic and Optometric Sciences*, 4, 47-49.
 25. Naser, M., & Shushtarian, S. M. M. (2020). Need for Visual Pathway Examination of the Patient Prior to Bone Marrow Transplantation. *Journal of Ophthalmology and Research*, 3, 75-78.
 26. Sanaie, S. J., Nematian, S., Shushtarian, M. M., & Khodaieani, A. R. (2014). Study of electrooculogram (EOG) abnormalities in patient with ocular toxoplasmosis. *Medical Science Journal of Islamic Azad University-Tehran Medical Branch*, 24, 33-36.
 27. Adhami-Moghadam, F., Talebi-Bidhendi, S., & Shushtarian, S. M. M. (2020). Retinal Screening of Workers Exposed to Mercury Vapor Using Electroretinography. *Journal of Ophthalmic and Optometric Sciences*, 4, 34-38.
 28. Shushtarian, S. M. M. (2020). Flash and Pattern Reversal Checkerboard Visual Evoked Potential Recording in Albinism Patients. *Journal of Ophthalmic and Optometric Sciences*, 4, 42-46.
 29. Fatemian, N., & Adhami-Moghadam, F. (2021). Seyed Mohammad Masoud Shushtarian (2021). Study of Visual Evoked Potentials in Patients Suffering from Exotropia. *Journal of Ophthalmic and Optometric Sciences*, 5.
 30. Shushtarian, S. M. M., & Pour Mazar, R. (2021). Far Distance Blurry Vision Following Rhinoplasty. *Journal of Ophthalmic and Optometric Sciences*, 5, 71-74.
 31. Shushtarian, S. M. M., Shojaei, A., & Tajik, F. (2018). Visual Pathway Disturbances in Rossai-Dorfman Disease: A Case Report. *Journal of Ophthalmic and Optometric Sciences*, 2, 24-26.
 32. Shushtarian, S. M. M., Shahmohammadi Dermani, F., &

- Pour Mazar, R. (2021). Blurred Vision in a Patient Suffering from Endometriosis and Epilepsy. *Journal of Ophthalmic and Optometric Sciences*, 5, 57-60.
33. Shushtarian, S. M. M. (2021). Dizziness and Nausea Feeling During Pattern Reversal Checkerboard Visual Evoked Potential Recording in A Multiple Sclerosis Patient. *Journal of Ophthalmic and Optometric Sciences*, 5, 44-47.
 34. Shushtarian, S. M. M., & Adhami-Moghadam, F. (2021). Low Vision in a Patient Due to Retinal Dystrophy upon Refractive Surgery. *Journal of Ophthalmic and Optometric Sciences*, 5.
 35. Adhami-Moghadam, P., Shushtarian, S. M. M., & Adhami-Moghadam, F. (2021). Retinal Screening of Coats Disease Using Electroretinography. *Journal of Ophthalmic and Optometric Sciences*, 5, 14-18.
 36. Ameli, S. (2023). Ali Panahi sharif, Sanaz saleh, seyed Mohammad Masoud Shushtarian, Ahmad Shojaei. Flash Visual Evoked Potential Recording in Patients with Brain Stroke. *Journal of Ophthalmology and Research*, 6, 40-44.
 37. Shushtarian, S. M. M., Jazayeri, S. Y., & Vafaei, A. (2023). Visual Evoked Potential Findings in Patients with Dyslexia. *Journal of Ophthalmic and Optometric Sciences*, 7.
 38. Shushtarian, S. M. M. (2022). Psychogenic Vomiting in a Child during Visual Evoked Potential Recording: A case Report. *Journal of Ophthalmic and Optometric Sciences*, 6, 33-36.
 39. Fatemian, N., Shushtarian, S. M. M., & Shojaei, A. (2022). Retinal Screening of Patients Suffering from Bardet-Biedl Syndrome Using Electroretinography. *Journal of Ophthalmic and Optometric Sciences*, 6(1).
 40. Pour Mazar, R. (2022). Retinal Screening of Patients Suffering from Bardet-Biedl Syndrome Using Electroretinography. *Journal of Ophthalmic and Optometric Sciences*, 6, 21-27.
 41. Shushtarian, S. M. (2009). Role of Myelin in Synchronization and Rhythmicity of Visual Impulses. In 4th European Conference of the International Federation for Medical and Biological Engineering (pp. 160-162).
 42. Shushtarian, S. M. M., & Hayti, Z. (2019). Probable Toxic Effect of Sodium Valproate on Retina Using Electroretinogram. *Journal of Ophthalmic and Optometric Sciences*, 3, 24-28.
 43. Shushtarian, S. M. M., & Mazar, R. P. (2022). A Rare Case of Visual Decline due to Intradiscal Ozone Therapy in a Patient with Spinal Stenosis. *Journal of Ophthalmic and Optometric Sciences*, 6, 27-29.
 44. Shushtarian, S. M. M., & Adhami-Moghadam, F. (2022). Retinal Screening of Coats Disease Using Electrooculography. *Journal of Ophthalmic and Optometric Sciences*, 6, 30-36.
 45. Shushtarian, S. M., Valiollahi, P., & Yahyabek, A. (2009). Effect of Distance between Monitor & Patient in Recording Visual Evoked Potential Using Pattern Reversal Checker Board Stimulation. In *World Congress on Medical Physics and Biomedical Engineering* (Vol. 25, pp. 861-863).
 46. Shushtarian, S. M. M., & Fatemian, N. (2021). Large Difference in Latency of Visual Evoked Potential P100 Peak in Case of Pattern and Flash Stimulation in a Multiple Sclerosis Patient. *Journal of Ophthalmic and Optometric Sciences*, 5, 73-76.
 47. Naser, M., & Shushtarian, S. M. M. (2020). Amplitude and Latency of Electroretinographical Peaks as a tool to predict the Extent of Retinal Degeneration in Retinitis Pigmentosa Patients. *Journal of Ophthalmology and Research*, 3, 71-74.
 48. Sarzaeim, F., Hashemzahi, M., Shushtarian, S. M. M., & Shojaei, A. (2022). Visual Evoked Potential Findings in Road Drilling Machine laborers. *Journal of Ophthalmology and Research*, 5, 43-47.
 49. Shushtarian, S. M. M., Shojaei, A., & Adhami-Moghadam, F. (2018). Visual Evoked Potentials Changes among Patients with Chronic Mustard Gas Exposure. *Journal of Ophthalmic and Optometric Sciences*, 2, 6-9.
 50. Sarzaeim, F., Abdolalizadeh, S., Shushtarian, S. M. M., & Shojaei, A. (2022). Visual Evoked Potential Findings in Patients using Anti-Seizure Medicine. *Journal of Ophthalmology and Research*, 5, 123-126.
 51. Renner, A. B., Kellner, U., Tillack, H., Kraus, H., & Foerster, M. H. (2005). Recording of both VEP and multifocal ERG for evaluation of unexplained visual loss: Electrophysiology in unexplained visual loss. *Documenta ophthalmologica*, 111, 149-157.
 52. Güngör, L., Güngör, İ., Öztürk, H. E., & Onar, M. K. (2011). Visual evoked potentials in guillain-barré syndrome. *Journal of Clinical Neurology*, 7, 34-39.
 53. Zgorzalewicz, M., Zielińska, M., & Kilarski, D. (2004). Auditory brainstem and visual evoked potentials in children and adolescents with Guillain-Barré syndrome [Brain stem auditory and visual evoked potentials in children and adolescents with Guillain-Barré syndrome]. *Neurol Neurochir Pol*, 38, S31-37.