

The Impact of Differences in Athletic Performance on Blood Indices in University Soccer Players

Ryunosuke Takahashi¹, Rena Murakami², Yukiko Kobayashi², & Takako Fujii^{3*}

¹The Institute of Health and Sports Science, Chuo University, Tokyo, Japan

²Graduate School of Life and Environmental Sciences, Kyoto Prefectural University, Kyoto, Japan

³Department of Sports and Medical Science, Kokushikan University, Tokyo, Japan

*Corresponding author: Takako Fujii, Department of Sports and Medical Science, Kokushikan University, Tokyo, Japan.

Submitted: 23 April 2025 Accepted: 30 April 2025 Published: 05 May 2025

doi DOI: <https://doi.org/10.63620/MKWJMM.2025.1019>

Citation: Fujii, T., Takahashi, R., Murakami, R., & Kobayashi, Y. (2025). The Impact of Differences in Athletic Performance on Blood Indices in University Soccer Players. *Word Jour of Molecule Medicine*, 2(3), 01-07.

Abstract

Background: Athletes need a diet that suits them to maintain and improve their performance. In recent years, dietary supplements have become more accessible and widely used by athletes, but there is concern that they may lead to reduced energy expenditure. In this study, physical characteristics of football players, blood composition and dietary surveys were conducted to investigate the effects of different athletic performance of athletes on iron nutritional status.

Methods: Dietary surveys, physical activity surveys, and measurements of physical characteristics were performed on 83 Japanese university soccer players.

Results: Body weight, muscle weight, reticulocytes, erythrocytes, hemoglobin, hematocrit and hepcidin were significantly higher in team A than in team B. Body fat mass and body fat percentage were significantly lower in team A than in team B. In terms of dietary content, potassium, iodine and niacin were significantly higher in team A than in team B.

Conclusion: There were no significant differences in dietary content. Although hepcidin levels were higher in team A than in team B, hemoglobin and ferritin levels were not reduced. Further studies are needed.

Keywords: Athlete, Nutrition, Diet, Iron, Hepcidin

Introduction

In 2010, the International Olympic Committee (IOC) stated that "a diet providing sufficient energy from a wide range of commonly available foods can meet the carbohydrate, protein, fat and micronutrient requirements for training and competition" [1]. Therefore, in our study "Dietary Self-Management in University Soccer Players: Are There Differences by Athletic Performance?" (Takahashi 2023), we reported that athletes with higher athletic performance had better dietary self-management skills [2]. Athletes with higher athletic ability consumed more iron than athletes with general athletic ability, and ferritin levels in blood indicators were higher in athletes with higher athletic ability. However, there was no difference in hemoglobin con-

centration.

Iron is a micronutrient found in very small amounts in the body, but it plays an important role in processes such as oxygen transport and energy metabolism. Iron is also an essential component of hemoglobin, which is required for oxygen transport in red blood cells. Therefore, if an athlete's body is deficient in iron, the oxygen-carrying capacity of active muscles is reduced, leading to a decrease in athletic performance. In other words, iron is also involved in oxygen transport to the muscles during exercise and plays an important role in energy production during exercise [3]. Iron deficiency therefore leads to reduced athletic performance in endurance sports [4].

In addition, iron metabolism disorders such as iron recycling from the spleen and macrophages and iron absorption in the duodenum via hepcidin are also associated with iron deficiency [3]. Exercise is known to increase the inflammatory cytokine interleukin-6 (IL-6), which is secreted 3 hours later and inhibits iron absorption. In other words, it has been suggested that athletes with high athletic ability who are highly active will secrete high levels of hepcidin and ferritin levels should be lower.

Therefore, focusing on hepcidin, this study investigated the effects of different sports on blood conditions.

Methods

Subjects

The subjects of this study were 83 people (ages: 19-22 years) who belong to a university soccer club. The purpose and content of the study were fully explained to the subjects, and they voluntarily agreed to participate in the study. In order to evaluate self-management ability based on athletic ability, the university athletes were divided into two groups: a high athletic ability group (Group A, $n = 31$), consisting of students whose members include many who will become professional soccer players from 2025, and a general athletic ability group (Group B, $n = 52$).

This study was approved by the ethics committee of the Kyoto Prefectural University (No. 309).

Measurement Items

Body Measurements

Height was measured with a height meter, and body weight, body fat percentage, and skeletal muscle weight were measured using the impedance method with an InBody 770 body composition measuring device (InBody Japan Inc.).

Blood Analysis

Blood samples were drawn from the antecubital vein after each subject had been seated quietly for at least 15 min. The samples were analyzed in a local commercial laboratory (Kyoto Microbiology Research Institute, Kyoto, Japan). Red blood cells (RBC), hemoglobin (Hb), hematocrit (Ht), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC), ferritin, hepcidin, unsaturated iron binding capacity (UIBC), ferritin, aspartate aminotransferase (AST), alanine aminotransferase (ALT), creatinine, platelets, C-reactive protein (CRP) and white blood cells (WBC) and haptoglobin were measured. Anemia was defined as an Hb level of <13 g/dL. Iron depletion was defined as a ferritin level of <20 μ g/L. Hemolysis was determined by the indirect bilirubin concentration, which is the total bilirubin minus the direct bilirubin concentration.

Dietary Survey

Participants were instructed to document their meals on two non-game days within the study period (July 2024) using the approximate quantity method and capture images with a smartphone. During photography, the participants were asked to include their student ID alongside the meal to aid in portion size determination. Nutrients and other intakes were analyzed from

the collected records and photographs using specialized nutrition calculation software (Eiyou Plus ver. 1.0; Kenpakusha, Tokyo).

Daily intakes and energy density-adjusted values were calculated for 22 items: energy; protein; fat; carbohydrate; potassium; calcium; magnesium; phosphorus; iron; zinc; copper; retinol activity equivalent; vitamin D, K, B₁, B₂, B₆, B₁₂, and C; alpha-tocopherol; folic acid; and salt equivalent. Additionally, data on the frequency and quantity of protein supplement consumption were gathered and incorporated into nutrient and other intake analyses. However, the nutrient content of supplements other than proteins, such as multivitamins, was not included in the analysis.

Physical Activity Survey

The physical activity questionnaire was administered over two days on the same day as the dietary questionnaire. Behavioural data were collected at approximately 5-minute intervals from awakening to bedtime using recording forms. Physical activity intensity was assessed using the Physical Activity Recording Standards for Health Promotion 2013,14 and the revised METs table for physical activity¹⁵. The intensity and duration of training at their club was determined by analysing videos of training sessions. The physical activity survey was completed over two days on the same day as the dietary survey. Behavioural data were collected at approximately 5-minute intervals from awakening to bedtime using recording forms. In this study, we therefore investigated the effects of differences in the athletes' sports on blood conditions, focusing on hepcidin.

Daily energy expenditure (kcal/day) was calculated using the formula¹⁶ was calculated using the formula: daily energy expenditure = body weight \times 1.05 \times Σ (exercise intensity \times time). The participants participated in club activities six days a week, with some undertaking independent training.

Statistical Analysis

Descriptive statistics were reported as mean (SD). Differences in means between the two groups were analysed using the t-test. The unpaired t-test was used to compare normally distributed data, whereas the Mann-Whitney U test was used to compare non-normally distributed data between groups. Pearson's correlation coefficient was used to assess correlation for normally distributed data, whereas Spearman's rank sum correlation coefficient was used for non-normally distributed data. Statistical analysis was performed using JMP Pro 18 (SAS Institute, Cary, NC, USA). P values <0.05 were considered to indicate statistical significance.

Results

The participants' physical characteristics are listed in Table 1. The height, weight, and muscle mass were significantly higher in Team A than in Team B. The body fat percentage was significantly lower in Team A than in Team B. There were no significant differences in the height, BMI, or body fat mass between the groups.

Table 1. Body composition		A (n=31)	B (n=52)
Age	year	20.6 (1.2)	20.4 (1.1)
High	cm	177.5 (5.5)	174.9 (6.4)
Weight	kg	70.6 (6.0)*	66.9 (5.9)
BMI	kg/m ²	22.4 (1.2)	21.9 (1.3)
Muscle mass	kg	60.4 (5.3)	56.0 (5.7)
Fat mass	kg	6.7 (2.1)*	7.7 (1.7)
Body fat percentage	%	9.3 (2.9)*	11.3 (2.8)
Activity	kcal	3919 (480)*	3625 (593)
mean (SD), *;p<0.05 vs B.			
BMI; Body mass index			

Table 2 shows the blood test results of the two groups. There were no anemic subjects in either group, and reticulocytes, RBCs, plasma iron, hemoglobin, hematocrit, and hepcidin were significantly higher than in Team B. Values for plasma iron, mean corpuscular volume, mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration, unsaturated iron binding capacity, ferritin, aspartate aminotransferase, alanine aminotransferase, creatinine, platelets, C-reactive protein, WBCs, and

haptoglobin did not differ significantly between groups. Table 3 shows a comparison of the nutritional intake of both groups. Group A had significantly higher levels of potassium, iodine, and niacin than Group B. The energy, protein fat, carbohydrate, calcium, magnesium, phosphorescence, iron, zinc, copper, iodine, vitamin A, D, E, K, B1, B2, B6, B12, folic acid, C, dietary fiber, and salt equivalent intake of the two groups did not differ significantly.

Table 2. Blood Indicators		A (n=31)	B (n=52)
Reticulocyte	/mL	17.5 (3.7)*	15.8 (4.0)
RBC	X10 ⁶ /μL	515 (24)*	500 (29)
MCV	fL	95.3 (3.1)	95.9 (2.9)
MCH	pg	29.7 (1.1)	29.9 (1.0)
MCHC	%	31.1 (0.6)	31.1 (0.7)
Hematocrit	%	49.0 (2.0)*	47.9 (2.4)
Plasma iron	μg/dL	108 (32)	106 (39)
Hemoglobin	g/dL	15.2 (0.6)*	14.9 (0.8)
UIBC	μg/dL	212 (49)	219 (52)
Ferritin	ng/mL	164 (54)	160 (65)
Hepcidin	μg/mL	20.0 (54.0)*	12.4 (9.7)
Haptoglobin	mg/dL	44.7 (29.2)	49.7 (40.6)
WBC	μL	5810 (1445)	5644 (1364)
Platelet count	× 10 ⁴ /μL	24.1 (4.3)	23.8 (3.8)
CRP	mg/dL	0.11 (0.15)	0.17 (0.42)
Creatinine	mg/dL	0.94 (0.11)	0.90 (0.09)
AST	U/l	22.2 (10.1)	22.0 (5.7)
ALT	U/l	13.9 (11.2)	12.4 (5.6)
mean (SD), *;p<0.05 vs B.			
MCV; Plasma iron, mean corpuscular volume. MCH; mean corpuscular hemoglobin. MCHC; mean corpuscular hemoglobin concentration. UIBC; unsaturated iron binding capacity. AST; aspartate aminotransferase. ALT; alanine aminotransferase. CRP; C-reactive protein,.			

Table 3. Dietary survey

		A (n=31)	B (n=52)
Energy	kcal	3243 (629)	3081 (898)
Protein	g	118 (25)	110 (32)
Fat	g	107 (32)	102 (37)
Cholesterol	mg	599 (210)	605 (256)
Carbohydrates	g	401 (93)	397 (152)
Potassium	mg	3787 (869)*	3290 (1196)
Calcium	mg	460 (149)	481 (238)
Magnesium	mg	389 (85)	351 (127)
Phosphorescence	mg	1629 (369)	1534 (448)
Iron	mg	12.0 (3.4)	11.4 (4.1)
Zine	mg	17.1 (4.4)	16.1 (5.5)
Copper	mg	1.9 (0.5)	1.8 (0.6)
Iodine	µg	3761 (2524)*	2012 (2471)
Vitamin A	µg	592 (244)	577 (293)
Vitamin D	µg	17.0 (12.0)	11.9 (14.2)
Vitamin E	mg	11.3 (4.7)	9.9 (4.2)
Vitamin K	µg	485 (273)	474 (312)
Vitamin B1	mg	2.2 (0.8)	1.9 (0.7)
Vitamin B2	mg	2.1 (0.8)	1.8 (0.7)
Niacin	mg	72 (19)*	56 (19)
Vitamin B6	mg	2.7 (0.7)	2.4 (0.9)
Vitamin B12	µg	9.7 (5.0)	8.4 (6.2)
Folic acid	µg	403 (141)	402 (194)
Vitamin C	mg	160 (109)	121 (82)
Dietary fiber	g	32 (8)	29 (10)
Salt equivalent	g	13 (3)	12 (3)

mean (SD), *;p<0.05 vs B.

Table 4. Correlation coefficient results for Team A

		<i>r</i>	<i>p</i>
Hepcidin	Reticulocyte	0.3610	0.0460*
Hepcidin	Feritin	0.4379	0.0137*
Hepcidin	WBC	0.4587	0.0095*
Hepcidin	CRP	0.1925	0.0021*
Hepcidin	AST	-0.4046	0.024*

*; p<0.05.

MCV; Plasma iron, mean corpuscular volume. MCH; mean corpuscular hemoglobin. MCHC; mean corpuscular hemoglobin concentration. UIBC; unsaturated iron binding capacity. AST; aspartate aminotransferase. ALT; alanine aminotransferase. CRP; C-reactive protein.

Screening for the Presence of Hepcidin and Related Substances

Regarding the correlation between hepcidin levels and body composition, blood indices, and dietary questionnaires, Pearson's correlation results showed significant positive correlations between reticulocytes, WBCs, creatinine, and ferritin in group

A ($p < 0.05$, $p < 0.04$, $p < 0.01$, and $p < 0.01$, respectively) (Table 4) whereas in group B, serum iron and ferritin levels showed a significant positive correlation ($p < 0.01$, $p < 0.01$, respectively) (Table 5).

Table 5. Correlation coefficient results for Team B

		r	p
Hepcidin	Plasma iron	0.2736	0.0497*
Hepcidin	UIBC	-0.313	0.0239*
Hepcidin	Ferritin	0.4503	0.0008*
Hepcidin	CRP	0.3271	0.0233*
Hepcidin	Body fat percentage	0.3498	0.0110*
Hepcidin	Fat mass	0.3696	0.0070*

*, $p < 0.05$.

MCV; Plasma iron, mean corpuscular volume. MCH; mean corpuscular hemoglobin. MCHC; mean corpuscular hemoglobin concentration. UIBC; unsaturated iron binding capacity. AST; aspartate aminotransferase. ALT; alanine aminotransferase. CRP; C-reactive protein,.

Spearman's analysis showed positive correlations for reticulocytes, WBCs, and ferritin in Team A ($p < 0.01$, $p < 0.01$, and $p < 0.01$, respectively); positive correlations for serum iron and ferritin in Team B ($p < 0.05$, $p < 0.01$, respectively); and positive

correlations for serum iron and ferritin in Team B ($p < 0.05$, $p < 0.01$, respectively). In the combined analysis of the two groups, body fat mass and percentage, WBCs, CRP level, and ferritin level were positively correlated with hepcidin levels (Table 6).

Table 6. Correlation coefficient results for soccer athlete (Team A and Team B)

		r	p
Hepcidin	UIBC	-0.3232	0.0029*
Hepcidin	Ferritin	0.3834	0.0003*
Hepcidin	WBC	0.2663	0.0149*
Hepcidin	CRP	0.2683	0.0142*
Hepcidin	AST	-0.2996	0.0059*
Hepcidin	Body fat percentage	0.2325	0.0344*
Hepcidin	Fat mass	0.2416	0.0278*

*, $p < 0.05$.

MCV; Plasma iron, mean corpuscular volume. MCH; mean corpuscular hemoglobin. MCHC; mean corpuscular hemoglobin concentration. UIBC; unsaturated iron binding capacity. AST; aspartate aminotransferase. ALT; alanine aminotransferase. CRP; C-reactive protein,.

Discussion

This study investigated the effects of iron nutritional status on different levels of competition in university soccer players. The results showed that erythrocyte and hemoglobin concentrations were significantly higher in team A than in team B. However, hepcidin was also significantly higher in team A than in team B.

Hemoglobin concentration was significantly higher in team A than in team B. This may be because Team A consumed more iron than Team B, but there was no significant difference compared to the dietary survey. In addition, the intake of vitamin C, a nutrient that enhances iron absorption, tended to be higher in team A than in team B ($p < 0.07$). These results suggest that iron absorption was increased, leading to the prevention of anemia. However, it should be noted that hemoglobin levels were within the normal range in both teams. In general, the recommended iron intake for athletes is 15 mg/day [5]. Both groups had ad-

equate hemoglobin concentrations, although they did not meet the 15 mg/day limit.

Ferritin concentrations were found in Team A and Team B. Mielgo Ayuso et al reported that 30-99 ng/ml is considered functional iron deficiency for screening for iron deficiency and a serum ferritin level of at least 100 ng/ml is required [6]. Both groups exceeded 100 ng/mL and are unlikely to have iron deficiency anemia.

On the other hand, hepcidin was significantly higher in Team A than in Team B. It has been reported that muscle glycogen decreases and hepcidin increases when energy intake remains inadequate [7]. When comparing the energy intake and activity of the athletes, the activity of both teams was higher than the energy intake: -700 kcal for team A and -500 kcal for team B. It is possible that the increased hepcidin in team A, which failed

to meet its energy intake requirements, could be due to insufficient energy intake. A correlation between hepcidin and WBC and CRP were also observed in the A team correlation; WBC and CRP reflect an inflammatory response. It is possible that Team A athletes have a chronic inflammatory state, especially as this was not a post-exercise blood sample.

The main sources of energy during exercise are carbohydrate and lipids. Carbohydrate in particular is known to be an important source of energy for athletes [8] and it has been reported that muscle glycogen is depleted at the point where exercise cannot be continued [9]. Williams & Rollo have highlighted the importance of having sufficient muscle glycogen stores for soccer player [10] and improved performance has been reported in players consuming high carbohydrate diets with dietary carbohydrate levels of ~8 g/kg BW [11, 12].

The latest nutritional guidelines for soccer recommend a daily carbohydrate intake of 5-7 g/kg body weight on days of moderate intensity training and 7-12 g/kg body weight on days of high intensity training and before matches [13]. The results of this study showed that Team A (5.7 g/kg body weight) and Team B (6.0 g/kg body weight). It has been reported that Asians rely more on carbohydrates for energy [14]. Based on the above, it is possible that increased carbohydrate intake may help both Team A and Team B to build a body that can withstand higher intensity and longer training sessions.

□

While the possibility of chronic inflammation was suggested in Team A, the hemoglobin concentration was significantly higher in Team A than in Team B, although the blood indices were in the normal range in both teams. Energy intake was also inadequate in both groups. It has been suggested that chronic low energy availability (LEA) may also affect biomarkers involved in erythrocyte turnover [15-19]. A study by Hennigar et al. in active men reported that a 55% reduction in EA over 28 days resulted in a decrease in hemoglobin concentration [20]. This reduction in hemoglobin (anemia), which does not respond to iron administration, suggests that it may be due to inhibition of red blood cell production by LEA [20]. □

When erythropoiesis is reduced as a result of energy deprivation, the number of erythroid progenitor cells in the bone marrow decreases, maintaining iron stores for vital functions. This response has been reported to underlie the development of iron-limiting anemias, including iron deficiency anemia and anemia due to chronic disease and inflammation [21]. Testosterone is also a potent stimulator of erythropoiesis [22-25]. However, the effects and mechanisms by which testosterone affects human erythropoiesis during severe energy deprivation remain unclear. Injection of testosterone into energy-restricted mice increased the incorporation of ⁵⁹Fe into newly formed erythrocytes, an indirect measure of erythropoiesis, compared to energy restriction alone [26].

Testosterone replacement has been reported to suppress hepcidin in healthy adults, thereby increasing iron utilisation and maintaining red blood cell production during energy deficiency [27]. It has also been suggested that testosterone administration may be an effective treatment for maintaining erythropoiesis during energy deficiency and may be particularly relevant for

certain types of anemia associated with chronic disease [28, 29]. Exercise is known to increase testosterone secretion. Although testosterone was not measured in this study, it is possible that the significantly higher hemoglobin concentration in team A, despite higher hepcidin concentration than in team B, was related to higher testosterone secretion due to higher activity levels.

It is known that obesity increases hepcidin levels and has been reported to be due to inflammation. In the present study, hepcidin levels were significantly higher in Team A than in Team B, even though Team A had less body fat than Team B. Nirengi et al. reported that hepcidin levels were higher in athletes with less body fat than in those with more body fat. Body fat is usually positively correlated with hepcidin concentration. However, there are few reports on the relationship between hepcidin and body fat in athletes. More research is needed on the relationship between hepcidin and body fat in athletes.

Conclusion

The present study investigated the effects of different levels of competition on blood indices, mainly iron nutritional status, in university soccer players. The results showed that erythrocyte and hemoglobin concentrations were significantly higher in team A than in team B, while hepcidin concentrations were significantly higher in team A than in team B.

Team A showed a positive correlation between hepcidin and reticulocyte, WBC and ferritin; Team B showed a positive correlation between hepcidin and plasma iron and ferritin.

There are few reports on indicators of exercise capacity differences in relation to hepcidin. More research is needed on the effects of differences in exercise capacity on hepcidin.

References

1. Maughan, R., & Shirreffs, S. (2010). IOC consensus statement on sports nutrition. *Journal of Sports Sciences*, 29(1), 3-4.
2. Takahashi, R., Ito, S., Hata, K., & Fujii, T. (2024). Dietary self-management in university soccer players: Are there differences by athletic performance? *Biomedical Journal of Scientific & Technical Research*, 54(4).
3. Fried, W., Goldwasser, E., Jacobson, L. O., & Plzak, L. F. (1957). Studies on erythropoiesis. III. Factors controlling erythropoietin production. *Proceedings of the Society for Experimental Biology and Medicine*, 94(1), 237-241. <https://doi.org/10.3181/00379727-94-22910>
4. Donati, R. M., Chapman, C. W., Warnecke, M. A., & Gallagher, N. I. (1964). Iron metabolism in acute starvation. *Proceedings of the Society for Experimental Biology and Medicine*, 117, 50-53. <https://doi.org/10.3181/00379727-117-29493>
5. Slimani, M., & Nikolaidis, P. T. (2019). Anthropometric and physiological characteristics of male soccer players according to their competitive level, playing position and age group: A systematic review. *Journal of Sports Medicine and Physical Fitness*, 59(1), 141-163. <https://doi.org/10.23736/S0022-4707.17.07950-6>
6. Abreu, R., Oliveira, C. B., Costa, J. A., Brito, J., & Teixeira, V. H. (2023). Effects of dietary supplements on athletic performance in elite soccer players: A systematic review. *Journal of the International Society of Sports Nutrition*, 20(1), 2236060. <https://doi.org/10.1080/15502783.2023.2236060>

7. Mielgo-Ayuso, J., Zourdos, M. C., Calleja-González, J., Córdova, A., Fernandez-Lázaro, D., & Caballero-García, A. (2018). Eleven weeks of iron supplementation does not maintain iron status for an entire competitive season in elite female volleyball players: A follow-up study. *Nutrients*, 10(10), 1526. <https://doi.org/10.3390/nu10101526>
8. Ishibashi, A., Maeda, N., Kojima, C., & Goto, K. (2022). Iron metabolism following twice a day endurance exercise in female long-distance runners. *Nutrients*, 14(9), 1907. <https://doi.org/10.3390/nu14091907>
9. Backhouse, S. H., Ali, A., Biddle, S. J., & Williams, C. (2007). Carbohydrate ingestion during prolonged high-intensity intermittent exercise: Impact on affect and perceived exertion. *Scandinavian Journal of Medicine & Science in Sports*, 17(5), 605-610. <https://doi.org/10.1111/j.1600-0838.2006.00613.x>
10. Balsom, P. D., Gaitanos, G. C., Söderlund, K., & Ekblom, B. (1999). High-intensity exercise and muscle glycogen availability in humans. *Acta Physiologica Scandinavica*, 165(4), 337-345. <https://doi.org/10.1046/j.1365-201x.1999.00517.x>
11. Williams, C., & Rollo, I. (2015). Carbohydrate nutrition and team sport performance. *Sports Medicine*, 45(Suppl 1), S13-S22. <https://doi.org/10.1007/s40279-015-0399-3>
12. Goedecke, J. H., White, N. J., Chicktay, W., Mahomed, H., Durandt, J., & Lambert, M. I. (2013). The effect of carbohydrate ingestion on performance during a simulated soccer match. *Nutrients*, 5(12), 5193-5204. <https://doi.org/10.3390/nu5125193>
13. Kingsley, M., Penas-Ruiz, C., Terry, C., & Russell, M. (2014). Effects of carbohydrate-hydration strategies on glucose metabolism, sprint performance and hydration during a soccer match simulation in recreational players. *Journal of Science and Medicine in Sport*, 17(2), 239-243. <https://doi.org/10.1016/j.jsams.2013.04.010>
14. Thomas, D. T., Erdman, K. A., & Burke, L. M. (2016). Position of the Academy of Nutrition and Dietetics, Dietitians of Canada, and the American College of Sports Medicine: Nutrition and athletic performance. *Journal of the Academy of Nutrition and Dietetics*, 116(3), 501-528. <https://doi.org/10.1016/j.jand.2015.12.006>
15. Sharma, S., Wilkens, L. R., Shen, L., & Kolonel, L. N. (2013). Dietary sources of five nutrients in ethnic groups represented in the Multiethnic Cohort. *British Journal of Nutrition*, 109(8), 1479-1489. <https://doi.org/10.1017/S0007114512003388>
16. Hennigar, S. R., Berryman, C. E., Harris, M. N., Karl, J. P., Lieberman, H. R., McClung, J. P., ... & Pasiakos, S. M. (2020). Testosterone administration during energy deficit suppresses hepcidin and increases iron availability for erythropoiesis. *The Journal of Clinical Endocrinology & Metabolism*, 105(4), e1316-e1321. <https://doi.org/10.1210/clinem/dgz316>
17. Pak, M., Lopez, M. A., Gabayan, V., Ganz, T., & Rivera, S. (2006). Suppression of hepcidin during anemia requires erythropoietic activity. *Blood*, 108(12), 3730-3735. <https://doi.org/10.1182/blood-2006-06-028787>
18. Fruhman, G. J. (1966). Effects of starvation and refeeding on erythropoiesis in mice. *Zeitschrift für Zellforschung und Mikroskopische Anatomie*, 75(1), 258-271. <https://doi.org/10.1007/BF00407159>
19. Gurney, C. W., & Fried, W. (1965). Further studies on the erythropoietic effect of androgens. *Journal of Laboratory and Clinical Medicine*, 65, 775-782.
20. Bachman, E., Trivison, T. G., Basaria, S., Davda, M. N., Guo, W., Li, M., Connor Westfall, J., Bae, H., Gordeuk, V., & Bhasin, S. (2014). Testosterone induces erythrocytosis via increased erythropoietin and suppressed hepcidin: Evidence for a new erythropoietin/hemoglobin set point. *Journal of Gerontology: Series A*, 69(6), 725-735. <https://doi.org/10.1093/geronol/glt154>
21. Suzuki, D., & Suzuki, Y. (2024). Identifying and analyzing low energy availability in athletes: The role of biomarkers and red blood cell turnover. *Nutrients*, 16(14), 2273. <https://doi.org/10.3390/nu16142273>
22. Ashby, D. R., Gale, D. P., Busbridge, M., Murphy, K. G., Duncan, N. D., Cairns, T. D., ... & Choi, P. (2009). Erythropoietin administration in humans causes a marked and prolonged reduction in circulating hepcidin. *haematologica*, 95(3), 505. <https://doi.org/10.3324/haematol.2009.013136>
23. Kautz, L., & Nemeth, E. (2014). Molecular liaisons between erythropoiesis and iron metabolism. *Blood*, 124(4), 479-482. <https://doi.org/10.1182/blood-2014-05-516252>
24. Zhang, L., Lang, H., Ran, L., Tian, G., Shen, H., Zhu, J., ... & Mi, M. (2022). Long-term high loading intensity of aerobic exercise improves skeletal muscle performance via the gut microbiota-testosterone axis. *Frontiers in Microbiology*, 13, 1049469. <https://doi.org/10.3389/fmicb.2022.1049469>
25. Nirengi, S., Fujibayashi, M., Furuno, S., Sukino, S., Suganuma, A., Kawaguchi, Y., ... & Sakane, N. (2021). The hepcidin-25/ferritin ratio is increased in university rugby players with lower fat mass. *Nutrients*, 13(9), 2993. <https://doi.org/10.3390/nu13092993>
26. Auguet, T., Aragonés, G., Berlanga, A., Martínez, S., Sabench, F., Binetti, J., ... & Richart, C. (2017). Hepcidin in morbidly obese women with non-alcoholic fatty liver disease. *PLoS One*, 12(10), e0187065. <https://doi.org/10.1371/journal.pone.0187065>
27. Andrews Guzmán, M., & Arredondo Olguín, M. (2014). Association between ferritin, high sensitivity C-reactive protein (hsCRP) and relative abundance of hepcidin mRNA with the risk of type 2 diabetes in obese subjects. *Nutrición Hospitalaria*, 30, 577-584.
28. Rodríguez-Mortera, R., Caccavello, R., Hermo, R., Garay-Sevilla, M. E., & Gugliucci, A. (2021). Higher hepcidin levels in adolescents with obesity are associated with metabolic syndrome dyslipidemia and visceral fat. *Antioxidants*, 10, 751. <https://doi.org/10.3390/antiox10050751>
29. Stoffel, N. U., El-Mallah, C., Herter-Aeberli, I., Bissani, N., Wehbe, N., Obeid, O., & Zimmermann, M. B. (2020). The effect of central obesity on inflammation, hepcidin, and iron metabolism in young women. *International Journal of Obesity*, 44, 1291-1300. <https://doi.org/10.1038/s41366-020-0564-6>