

Dynamic Sets S1et and Some of their Applications in Physics

Oleksandr Danilishyn and Illia Danilishyn*

Sumy State University, Ukraine

*Corresponding author: Illia Danilishyn, Sumy State University, Vulytsya Mykoly Sumtsova, Sumy Oblast, 40000, Ukraine.

Submitted: 11 September 2023 Accepted: 19 September 2023 Published: 25 September 2023

Citation: Oleksandr Danilishyn, Illia Danilishyn (2023) Dynamic Sets S1et and Some of Their Applications in Physics. Sci Set J of Physics 2(3), 01-11.

Abstract

The article aims to create new constructive hierarchical mathematical objects for new technologies, particularly for a fundamentally new type of neural network with parallel computing and not the usual parallel computing through sequential computing.

Keywords: Dynamic Set S1et, S1et-Elements, Capacity S1et, S1et-Sets in Themselves, S1et-Elements in Themselves, Sit-Elements, Capacity.

S¹et-Elements

Here, the axiom of regularity (A8) is removed from the axioms of set theory, so we naturally obtain the possibility of using singularities in the form of S1et - sets in themselves, S1et - elements in themselves, which is exactly what we need for new mathematical models for describing complex processes, in particular in physics [1]. Instead of the axiom of regularity, we introduce the following axioms: Axiom R1. $\forall B (St_{C_0B}^C = B)$. Axiom R2. $\forall B (\exists B^{-1})$.

Definition 1

The expression

$$C_1 S^1 t_{B_1}^{A_k}, j, k=1,2 (*)$$

where A_1, A_2 is contained into B_1 , D_1, D_2 is expelled from C_1 and for sstructure (*) is performed the next operation- multiplication:

$$C_1 S^1 t_{B_1}^{A_1} * C_2 S^1 t_{B_1}^{A_2} =_{D_1 \cup D_2} C_1 S^1 t_{B_1}^{A_1 \cup A_2} (*_1),$$

then we shall call S1et- elements, in case $A_1, A_2, B_1, D_1, D_2, C_1$ are sets we shall call (*) the dynamical hierarchical set S1et. $A_1, A_2, B_1, D_1, D_2, C_1$ -are any, in particular, $A_i, i=1, 2$, may be actions in the right direction, actions with the right goal (action with the so-called target weights, any actions [2]).

Definition 2

$C_D S^1 t_B^A$ is called an ordered S1et- element, if some or any elements from A, B, C, D may be by ordered elements.

where some or any elements may be by ordered elements.

S¹et- elements can be elements of a group by multiplication (*1).

S¹et-Capacity in Itself

Definition 3

The S¹et -capacity A in itself and from itself of the null type is the capacity containing itself as an element and expelling oneself out of oneself simultaneously:

$$A S^1 t_A^A. \text{ Denote } S^1_0 f_A.$$

Definition 4

The S¹et -capacity in itself A and from itself B of the first type is the capacity containing A itself as an element and expelling B oneself out of oneself simultaneously:

$$B S^1 t_A^A. \text{ Denote } S^1_1 f_B^A.$$

Definition 5

The S¹et ¹-capacity of the second type is the capacity containing B into A and expelling B oneself out of oneself simultaneously:

$$B S^1 t_A^B. \text{ Denote } S^1_2 f_B^A.$$

Definition 6

The S¹et ¹-capacity of the third type is the capacity containing B itself as an element and the displacement of B from A simultaneously:

$$B S^1 t_B^B. \text{ Denote } S^1_3 f_B^A.$$

Definition 7

S¹et -capacity A in itself of the fourth type is the capacity that contains the program that allows it to be generated and it to be degenerated simultaneously. Let's denote

$$S_4^{1et} fA.$$

Definition 8

S¹et -capacity A in itself of the fifth type contains itself in part and expelling oneself in part or contains a program that allows it to be generated in part and it to be degenerated in part, or both simultaneously. Let us denote

$$S_5^{1et} fA.$$

Connection of S¹et-Elements with S¹et-Capacity in Itself

Consider a fifth type of self- capacity. For example, based on

$$S_5^{1et} fA.$$

where $A=(a_1, a_2, \dots, a_n)$ it is possible to consider S¹et -capacity

in itself

$$S_5^{1et} fA.$$

with m elements from A, at $m < n$, which is formed by the form: $W_{m_n} = (m, (n, 1))$ (1)

that is, only m elements are located in the structure

$$S_5^{1et} fA.$$

S¹et -capacity in itself of the fifth type can be formed for any other structure, not necessarily S¹et, only through the obligatory reduction in the number of elements in the structure. In particular, using the form.

$$w_{m_1 \dots m_n} = (m_1, (m_2, (\dots (m_n, 1) \dots))) \quad (2)$$

Structures more complex than $S_5^{1et} fA$. can be introduced.

Mathematics S¹et Itself

1. Similarly, for the simultaneous execution of various operators:

$$\begin{matrix} F_0 C \\ F_1 D \end{matrix} S_1^{F_2 A} t_{F_3 B}^A$$

where F_0, F_1, F_2, F_3 are operators.

2. Similarly, for the simultaneous execution of various operators:

$$S_5^{1et} fA.$$

3. $j=0, 1, 2, 3, 4, 5$, where $\{F\} = (F_0, F_1, F_2, F_3, F_4, F_5)$ are operators.

Operations are taking place:

$$4. \quad \begin{matrix} B \\ D \end{matrix} S_1^{t_A^A} t_{B-B}^A = \begin{pmatrix} B \\ B \end{pmatrix} S_1^{t_A^A} t_{B-B}^A, \quad \mu(B_S^1 t_A^A) = \begin{pmatrix} \mu^S(B_S^1 t_A^A) \\ \mu(A) + 2\mu(B) - \mu(D - A) \end{pmatrix}$$

$$5. \quad \begin{matrix} B \\ A \end{matrix} S_1^{t_A^A} t_{B-B}^A = \begin{pmatrix} B \\ A \end{pmatrix} S_1^{t_A^A} t_{B-B}^A, \quad \mu(B_S^1 t_A^A) = \begin{pmatrix} \mu^{SS}(B_S^1 t_A^A) \\ 2\mu(B) - \mu(A) + \mu(A - B) \end{pmatrix}$$

$$6. \quad \begin{matrix} B \\ A \end{matrix} S_1^{t_A^A} t_{B-B}^A = \begin{pmatrix} B \\ A \end{pmatrix} S_1^{t_A^A} t_{B-B}^A, \quad \mu(B_S^1 t_A^A) = \begin{pmatrix} \mu^{SS}(B_S^1 t_A^A) \\ 2\mu(B) + \mu(A) - \mu(A - B) \end{pmatrix}$$

$$7. \quad \begin{matrix} B \\ Q \end{matrix} S_1^{t_A^A} t_{B-B}^A = \begin{pmatrix} S_1^{et} fB \\ Q-B \end{pmatrix} S_1^{t_A^A} t_{B-B}^A, \quad \mu(B_S^1 t_A^A) = \begin{pmatrix} \mu^{SS}(S_1^{et} fB) \\ 2\mu(B) + \mu(A - B) - \mu(Q - B) \end{pmatrix}$$

$$8. \quad \begin{matrix} B \\ Q \end{matrix} S_1^{t_A^A} t_{B-B}^A = \begin{pmatrix} B \\ Q \end{pmatrix} S_1^{t_A^A} t_{B-B}^A, \quad \mu(B_S^1 t_A^A) = \begin{pmatrix} \mu^{SS}(B_S^1 t_A^A) \\ 2\mu(B) + \mu(A - B) - \mu(Q) \end{pmatrix}$$

$$9. \quad \begin{matrix} B \\ Q \end{matrix} S_1^{t_A^A} t_{B-B}^A = \begin{pmatrix} B \\ Q \end{pmatrix} S_1^{t_A^A} t_{B-B}^A, \quad \mu(B_S^1 t_A^A) = \begin{pmatrix} \mu^{SS}(B_S^1 t_A^A) \\ 2\mu(B) + \mu(A) - \mu(Q - B) \end{pmatrix}$$

$$10. \quad \begin{matrix} R \\ Q \end{matrix} S_1^{t_A^A} t_{B-B}^A = \begin{pmatrix} S_1^{et} fR \\ Q-B \end{pmatrix} S_1^{t_A^A} t_{B-B}^A, \quad \mu(B_S^1 t_A^A) = \begin{pmatrix} \mu^{SS}(S_1^{et} fR) \\ \mu(B) + \mu(A - B) + \mu(R) - \mu(Q - B) \end{pmatrix}$$

The concepts of S¹et – force: $\begin{matrix} F_3 \\ F_4 \end{matrix} S_1^{t_{F_2}^{F_1}} t_{F_2}^{F_1}$ -the containment of force

F_1 into force F_2 and the displacement of force F_4 from force

F_3 simultaneously, S¹et – energy: $\begin{matrix} E_3 \\ E_4 \end{matrix} S_1^{t_{E_2}^{E_1}} t_{E_2}^{E_1}$ -the containment of

energy E_1 into energy E_2 and the displacement of energy E_4 from energy E_3 simultaneously.

Consider the concepts of S¹et -capacity in itself of physical objects A, B. Similar to the concepts of publication: the S¹et -capacity in itself of the first type is the capacity containing A itself as an element and expelling B oneself out of oneself si

multaneously: $\begin{matrix} B \\ B \end{matrix} S_1^{t_A^A} t_{A-A}^A$, S¹et -capacity in itself of the third type

contains itself in part and expelling oneself in part or contains a program that allows it to be generated and it to be degenerated

simultaneously partially, or both: $S_5^{1et} fA$, $S_5^{1et} fB$.

By analogy, for

$$S_0^{1et} fA, S_2^{1et} fB, S_3^{1et} fB, S_4^{1et} fA.$$

Also, you can consider these types of S¹et -capacity in itself for other objects.

For example: $S_i^{1et} f$ operator A, $S_i^{1et} f$ action B, $S_i^{1et} f$ made Q

$i=0, 1, 2, 3, 4, 5$ and etc.

Remark. The concept of elements of physics S¹et is introduced for energy space. The corresponding concept of elements of chemistry S¹et is introduced accordingly.

Dynamical S¹et-Elements

Definition 9

The process of the containment of A(t) into B(t) and the displacement of D(t) from C(t) at time to simultaneously we shall call dynamical S¹et – element. Let's denote

$$\begin{matrix} C(t) \\ D(t) \end{matrix} S_1^{t(t)} t_{B(t)}^{A(t)}.$$

Definition 10

$\begin{matrix} C(t) \\ D(t) \end{matrix} S_1^{t(t)} t_{B(t)}^{A(t)}$ with ordered elements $\vec{A}(t)$ and $\vec{D}(t)$ is called

an ordered dynamical S¹et – element.

It is allowed to multiply dynamical S1et-elements:

$$_{D_1(t)}^{C_1(t)}S^1t(t)_{B_1(t)}^{A_1(t)} * _{D_2(t)}^{C_2(t)}S^1t(t)_{B_2(t)}^{A_2(t)} = _{D_1(t) \cup D_2(t)}^{C_1(t)}S^1t(t)_{B_1(t)}^{A_1(t) \cup A_2(t)} (*_2)$$

where some or any elements may be by ordered elements.

Dynamical S1et – elements can be elements of a group by multiplication (*2).

Dynamical S1et -Capacity in Itself

Definition 11

The dynamical S1et -capacity A(t) in itself and from itself of the null type is the process of a containment itself as an element and expelling oneself out of oneself at time t simultaneously:

$$_{A(t)}^{A(t)}S^1t(t)_{A(t)}^{A(t)}. \text{ Denote } S^{1et}_0(t)fA(t).$$

Definition 12

The dynamical S2et -capacity in itself A(t) and from itself B(t) of the first type is the process of a containment A(t) in itself as an element and expelling B(t) oneself out of oneself at time t simultaneously:

$$_{B(t)}^{B(t)}S^1t(t)_{A(t)}^{A(t)}. \text{ Denote } S^{1et}_1(t)f_{B(t)}^{A(t)}.$$

Definition 13

The dynamical S1et 1-capacity of the second type is the process of putting B(t) into A(t) and expelling B(t) oneself out of oneself at time t simultaneously:

$$_{B(t)}^{B(t)}S^1t(t)_{A(t)}^{B(t)}. \text{ Denote } S^{1et}_2(t)f_{B(t)}^{A(t)}.$$

Definition 14

Dynamical S1et1-capacity of the third type is the process of a containment of B(t) in itself as an element and the displacement of B(t) from A(t) at time to simultaneously:

$$_{B(t)}^{A(t)}S^1t(t)_{B(t)}^{B(t)}. \text{ Denote } S^{1et}_3(t)f_{B(t)}^{A(t)}.$$

Definition 15

Dynamical S1et -capacity A(t) in itself of the fourth type is the process of a containment of the program that allows it to be generated and it to be degenerated at time t simultaneously through the structure S1et. Let's denote

$$S^{1et}_4(t)fA(t).$$

Definition 16

Dynamical S1et -capacity A(t) in itself of the fifth type is the process of a containment of itself in part and expelling oneself in part or process of a containment of the program that allows it to be generated in part and it to be degenerated in part at time t through the structure S1et, or both simultaneously. Let us denote

$$S^{1et}_5(t)fA(t).$$

Consider dynamical S1et -capacity A(t) in itself of the fifth type:

$$S^{1et}_5(t)fA(t).$$

For $A(t)=(a_1(t), a_2(t), \dots, a_n(t))$ it is possible to consider the dynamical S1et -capacity A(t) in itself of the fifth type:

$$S^{1et}_5(t)fA(t).$$

with m elements and from $\{a(t)\}$, at $m \leq n$, which is process to be formed by the form (1), that is, only m elements from A(t) are located in the structure

$$_{D(t)}^{C(t)}S^1t(t)_{B(t)}^{A(t)}.$$

The same for $D(t)=(d_1(t), d_2(t), \dots, d_n(t))$ in it. Dynamical S1et -capacity in itself of the fifth type can be formed for any other structure, not necessarily S1et, only through the obligatory reduction in the number of elements in the structure. In particular, using the form (2). Structures more complex than

$$S^{1et}_5(t)fA(t). \text{ can be introduced.}$$

Dynamical Mathematics S1et-Itself

1. Similarly, for the simultaneous execution of various operators:

$$_{F_1(t)D(t)}^{F_0(t)C(t)}S^1t(t)_{F_3(t)B(t)}^{F_2(t)A(t)}, \text{ where } F_0(t), F_1(t), F_2(t), F_3(t)$$

2. Similarly, for the simultaneous execution of various operators:

$$S^{1et}_j(t)fF(t)A(t), j=0,4,5, \text{ and } S^{1et}_k(t)f_{B(t)}^{A(t)}, k=1,2,3,$$

where $\{F(t)\} = (F_0(t), F_1(t), F_2(t), F_3(t))$ are operators.

$$_{D(t)}^{B(t)}S^1t(t)_{B(t)}^{A(t)} = \left(\begin{matrix} B(t)S^1t(t)_{B(t)}^{A(t)} * \\ B(t)S^1t(t)_{B(t)}^{A(t)} \end{matrix} \right),$$

3.

$$\mu_{D(t)}^{B(t)}S^1t(t)_{B(t)}^{A(t)} = \left(\begin{matrix} \mu^s(B(t)S^1t(t)_{B(t)}^{A(t)}) \\ \mu(A(t)) + 2\mu(B(t)) - \mu(D(t) - A(t)) \end{matrix} \right)$$

$$_{A(t)}^{B(t)}S^1t(t)_{B(t)}^{A(t)} = \left(\begin{matrix} B(t)S^1t(t)_{B(t)}^{B(t)} * \\ B(t)S^1t(t)_{B(t)}^{A(t)-B(t)} \end{matrix} \right),$$

$$\mu_{A(t)}^{B(t)}S^1t(t)_{B(t)}^{A(t)} = \left(\begin{matrix} \mu^{ss}(B(t)S^1t(t)_{B(t)}^{B(t)}) \\ 2\mu(B(t)) + \mu(A(t) - B(t)) - \mu(A(t)) \end{matrix} \right)$$

$$_{A(t)}^{B(t)}S^1t(t)_{B(t)}^{A(t)} = \left(\begin{matrix} B(t)S^1t(t)_{B(t)}^{A(t)} \\ B(t)S^1t(t)_{B(t)}^{A(t)} \end{matrix} \right),$$

$$\mu_{A(t)}^{B(t)}S^1t(t)_{B(t)}^{A(t)} = \left(\begin{matrix} \mu^{ss}(B(t)S^1t(t)_{B(t)}^{A(t)}) \\ 2\mu(B(t)) + \mu(A(t)) - \mu(A(t) - B(t)) \end{matrix} \right)$$

$$_{Q(t)}^{B(t)}S^1t(t)_{B(t)}^{A(t)} = \left(\begin{matrix} S^{1et}_0(t)f(t)B(t) * \\ B(t)S^1t(t)_{B(t)}^{A(t)-B(t)} \end{matrix} \right),$$

$$\mu_{Q(t)}^{B(t)}S^1t(t)_{B(t)}^{A(t)} = \left(\begin{matrix} \mu^{ss}(S^{1et}_0(t)f(t)B(t)) \\ 2\mu(B(t)) + \mu(A(t) - B(t)) - \mu(Q(t) - B(t)) \end{matrix} \right)$$

$$_{Q(t)}^{B(t)}S^1t(t)_{B(t)}^{A(t)} = \left(\begin{matrix} B(t)S^1t(t)_{B(t)}^{B(t)} * \\ B(t)S^1t(t)_{B(t)}^{A(t)-B(t)} \end{matrix} \right),$$

$$\mu_{Q(t)}^{B(t)}S^1t(t)_{B(t)}^{A(t)} = \left(\begin{matrix} \mu^{ss}(B(t)S^1t(t)_{B(t)}^{B(t)}) \\ 2\mu(B(t)) + \mu(A(t) - B(t)) - \mu(Q(t)) \end{matrix} \right)$$

$$10. \quad \begin{matrix} B(t) \\ Q(t) \end{matrix} S^1 t(t)_{B(t)}^{A(t)} = \left(\begin{matrix} B(t) \\ B(t) \end{matrix} S^1 t(t)_{B(t)}^{A(t)} * \right),$$

$$\mu_{Q(t)}^{B(t)} S^1 t(t)_{B(t)}^{A(t)} = \left(\begin{matrix} \mu^{ss} \left(\begin{matrix} B(t) \\ B(t) \end{matrix} S^1 t(t)_{B(t)}^{A(t)} * \right) \\ 2\mu(B(t)) + \mu(A(t)) - \mu(Q(t) - (B(t))) \end{matrix} \right)$$

$$11. \quad \begin{matrix} R(t) \\ Q(t) \end{matrix} S^1 t(t)_{B(t)}^{A(t)} = \left(\begin{matrix} S^{1et}_3 f(t)_{R(t)}^{B(t)} * \\ R(t) \end{matrix} S^1 t(t)_{B(t)}^{A(t)-B(t)} \right),$$

$$\mu_{Q(t)}^{R(t)} S^1 t(t)_{B(t)}^{A(t)} = \left(\begin{matrix} \mu^{ss} \left(S^{1et}_3 f(t)_{R(t)}^{B(t)} * \right) \\ \mu(B(t)) + \mu(A(t) - B(t)) + \mu(R(t)) - \mu(Q(t) - B(t)) \end{matrix} \right)$$

The concepts of dynamical S²et –force:

$$\begin{matrix} F_3(t) \\ F_4(t) \end{matrix} S^1 t(t)_{F_2(t)}^{F_1(t)}$$

the containment of force $F_1(t)$ into force $F_2(t)$ and the displacement of force $F_4(t)$ from force $F_3(t)$ at time to simultaneously, dynamical S¹et – energy:

$$\begin{matrix} E_3(t) \\ E_4(t) \end{matrix} S^1 t(t)_{E_2(t)}^{E_1(t)}$$

the containment of energy $E_1(t)$ into energy $E_2(t)$ and the displacement of energy $E_4(t)$ from energy $E_3(t)$ at time to simultaneously.

Consider the concepts of dynamical S¹et -capacity in itself of physical objects $A(t)$, $B(t)$. Similar to the concepts of publication: the dynamical S¹et -capacity in itself of the null type is the dynamical capacity containing itself as an element and expelling oneself out of oneself at time t simultaneously:

$$S^{1et}_0(t) fA(t) = \begin{matrix} A(t) \\ A(t) \end{matrix} S^1 t(t)_{A(t)}^{A(t)},$$

dynamical S¹et -capacity in itself of the fifth type contains itself in part and expelling oneself in part or contains a program that allows it to be generated and it to be degenerated at time t simultaneously partially, or both:

$$S^{1et}_k(t) f_{B(t)}^{A(t)} \quad S^{1et}_k(t) f_{B(t)}^{A(t)},$$

By analogy, for

$$S^{1et}_k(t) f_{B(t)}^{A(t)} \quad S^{1et}_k(t) f_{B(t)}^{A(t)}, \quad S^{1et}_3(t) f_{B(t)}^{A(t)}, S^{1et}_4(t) fA(t).$$

Also, you can consider these types of dynamical S¹et -capacity in itself for other objects. For example: $S^{1et}_i(t) f$

(t)f operator $A(t)$, $S^{1et}_i(t) f$

(t)f action $B(t)$, $S^{1et}_i(t) f$

made $Q(t)$ $i=0, 1, 2, 3, 4, 5$ and etc.

Remark. The concept of elements of physics dynamical S¹et is introduced for energy space. The corresponding concept of elements of chemistry dynamical S¹et is introduced accordingly.

S¹et-Elements for Continual Sets

Here we consider some continual S¹et -elements and continual self-capacity in itself as an element.

Definition 17

The containment of A into B and the displacement of D from C simultaneously, where A, B, D, C - sets of continual elements we shall call continual S¹et – element. Let's denote ${}^C_S S^1 t_B^A$.

Definition 18

${}^C_{\vec{D}} S^1 t_{\vec{B}}^{\vec{A}}$ with ordered elements \vec{A} and \vec{B} , where A, B, D, C - sets of continual elements, is called an ordered S¹et – element.

It is allowed to multiply continual S¹et – elements:

$${}^{C_1}_{D_1} S^1 t_{B_1}^{A_1} * {}^{C_2}_{D_2} S^1 t_{B_2}^{A_2} = {}^{C_1}_{D_1 \cup D_2} S^1 t_{B_1}^{A_1 \cup A_2} (*_3),$$

where some or any elements may be by ordered elements.

Continual S¹et – elements can be elements of a group by multiplication (*₃).

S¹et -Capacity in Itself for Continual Sets

Definition 19

The continual S¹et -capacity A in itself and from itself of the null type is the capacity containing itself as an element and expelling oneself out of oneself simultaneously, where A - set of continual elements: ${}^A_A S^1 t_A^A$. Denote $S^{1et}_0 fA$.

Definition 20

The ordered continual S¹et -capacity \vec{A} in itself and from itself of the null type is the capacity containing itself as an element and expelling oneself out of oneself simultaneously, where \vec{A} ordered set of continual elements: ${}^{\vec{A}}_{\vec{A}} S^1 t_{\vec{A}}^{\vec{A}}$. Denote $S^{1et}_0 f\vec{A}$

Definition 21

The continual S¹et -capacity in itself A and from itself B of the first type is the capacity containing A itself as an element and expelling B oneself out of oneself simultaneously, where A, B - sets of continual elements: ${}^B_B S^1 t_A^B$. Denote $S^{1et}_1 f_B^A$.

Definition 22

The continual S¹et 1-capacity of the second type is the capacity containing B into A and expelling B oneself out of oneself simultaneously, where A, B - sets of continual elements:

$${}^B_B S^1 t_A^B. \text{ Denote } S^{1et}_2 f_B^A.$$

Definition 23

The continual S¹et 1-capacity of the third type is the capacity containing B itself as an element and the displacement of B from A simultaneously, where A, B - sets of continual elements:

$${}^A_B S^1 t_B^B. \text{ Denote } S^{1et}_3 f_B^A.$$

Definition 24

The continual S¹et -capacity A in itself of the fourth type is the capacity that contains the program that allows it to be generated and it to be degenerated simultaneously, where A - set of continual elements. Let's denote $S^{1et}_4 fA$.

Definition 25

The continual S¹et -capacity A in itself of the fifth type contains itself in part and expelling oneself in part or contains a program that allows it to be generated in part and it to be degenerated in part simultaneously, or both, where A - set of continual elements. Let us denote $S^{1et}_5 fA$.

Definition 26

The ordered continual S¹et -capacity in itself \vec{A} and from itself B of the first type is the capacity containing \vec{A} itself as an element and expelling B oneself out of oneself simultaneously, where \vec{A} - ordered set of continual elements, B- set of continual elements: ${}^B_S S^1 t_{\vec{A}}^{\vec{A}}$. Denote $S^1_{11}{}^{et} f_{\vec{B}}^{\vec{A}}$.

Definition 27

The ordered continual S¹et 1-capacity in itself A and from itself \vec{B} of the first type is the capacity containing A itself as an element and expelling \vec{B} oneself out of oneself simultaneously, where \vec{B} -ordered set of continual elements, A- set of continual elements: ${}^{\vec{B}}_S S^1 t_A^{\vec{A}}$. Denote $S^1_{11}{}^{et} f_{\vec{B}}^{\vec{A}}$.

Definition 28

The ordered continual S¹et 2-capacity in itself \vec{A} and from itself \vec{B} of the first type is the capacity containing \vec{A} itself as an element and expelling \vec{B} oneself out of oneself simultaneously, where \vec{A}, \vec{B} -ordered sets of continual elements: ${}^{\vec{B}}_S S^1 t_{\vec{A}}^{\vec{A}}$. Denote $S^1_{11}{}^{et} f_{\vec{B}}^{\vec{A}}$.

Definition 29

The continual S¹et¹-capacity of the second type is the capacity containing B into \vec{A} and expelling B oneself out of oneself simultaneously, where \vec{A} - ordered set of continual elements, B- set of continual elements:

Definition 30

The continual S¹et 2-capacity of the second type is the capacity containing \vec{B} into A and expelling \vec{B} oneself out of oneself simultaneously, where \vec{B} -ordered set of continual elements, A- set of continual elements: ${}^{\vec{B}}_S S^1 t_A^{\vec{A}}$. Denote $S^1_{22}{}^{et} f_{\vec{B}}^{\vec{A}}$.

Definition 31

The continual S¹et 3-capacity of the second type is the capacity containing B into A and expelling \vec{B} oneself out of oneself simultaneously, where A, B -ordered sets of continual elements:

$${}^{\vec{B}}_S S^1 t_A^{\vec{A}}. \text{ Denote } S^1_{22}{}^{et} f_{\vec{B}}^{\vec{A}}.$$

Definition 32

The continual S¹et 1-capacity of the third type is the capacity containing B itself as an element and the displacement of B from A simultaneously, where \vec{A} - ordered set of continual elements, B- set of continual elements:

$${}^A_S S^1 t_{\vec{B}}^{\vec{B}}. \text{ Denote } S^1_{33}{}^{et} f_{\vec{B}}^{\vec{A}}.$$

Definition 33

The continual S¹et 2-capacity of the third type is the capacity containing \vec{B} itself as an element and the displacement of \vec{B} from A simultaneously, where B ordered set of continual elements, A- set of continual elements:

$${}^A_S S^1 t_{\vec{B}}^{\vec{B}}. \text{ Denote } S^1_{33}{}^{et} f_{\vec{B}}^{\vec{A}}.$$

Definition 34

The continual S¹et 3-capacity of the third type is the capacity containing \vec{B} itself as an element and the displacement of \vec{B} from A simultaneously, where A, B- ordered sets of continual elements:

$${}^A_S S^1 t_{\vec{B}}^{\vec{B}}. \text{ Denote } S^1_{33}{}^{et} f_{\vec{B}}^{\vec{A}}.$$

Definition 35

The ordered continual S¹et -capacity \vec{A} in itself of the fourth type is the capacity that contains the program that allows it to be generated and it to be degenerated simultaneously, where \vec{A} - set of continual elements. Let's denote: $S^1_{44}{}^{et} f_{\vec{A}}$.

Definition 36

The ordered continual S¹et -capacity \vec{A} in itself of the fifth type contains itself in part and expelling oneself in part or contains a program that allows it to be generated in part and it to be degenerated in part simultaneously, or both simultaneously, where \vec{A} - ordered set of continual elements. Let us denote

$$S^1_{55}{}^{et} f_{\vec{A}}.$$

Also we consider next elements

$$S^2_{00}{}^{et} f_{\vec{A}} \uparrow \downarrow \vec{A}, S^2_{11}{}^{et} f_{\vec{B}} \uparrow \downarrow \vec{A}, S^2_{22}{}^{et} f_{\vec{B}} \uparrow \downarrow \vec{A}, S^2_{33}{}^{et} f_{\vec{B}} \uparrow \downarrow \vec{A}, S^2_{44}{}^{et} f_{\vec{B}} \uparrow \downarrow \vec{A}, S^2_{55}{}^{et} f_{\vec{B}} \uparrow \downarrow \vec{A} \text{ etc.}$$

Connection of S¹et-Elements with Self-capacity in Itself as an Element

Consider a fifth type of continual self- capacity in itself as an element.

For example,

$S^1_{55}{}^{et} f_{\vec{A}}$, where $A=(a_1, a_2, \dots, a_n)$, i.e. a_i - continual elements, $i=1, 2, \dots, n$. It's possible to consider the continual self- capacity in itself as an element $S^1_{55}{}^{et} f_{\vec{A}}$, with m continual elements $S^1_{55}{}^{et} f_{\vec{A}}$, at $m < n$, which is formed by the form (1), that is, only m continual elements are located in the structure

$$S^1_{55}{}^{et} f_{\vec{A}},$$

Ccontinual self-capacity in itself as an element of the fifth type can be formed for any other structure, not necessarily S¹et, only through the obligatory reduction in the number of continual elements in the structure. In particular, using the form (2).

The same for $S^1_{55}{}^{et} f_{\vec{A}}$, Structures more complex than $S^1_{55}{}^{et} f_{\vec{A}}$, can be introduced.

Mathematics Itself for Continual S¹et -Elements

1. Simultaneous addition of a sets A, B, C, D with continual elements is realized by ${}^C_D S^1 t_{B \cup A}^{\vec{A}}$, where A, B, C, D may be ordered sets of continual elements.
2. Let's introduce operator to transform capacity to self-consistency in itself as an element: $Q_i S^1 \text{et}(A)$ transforms A to $S^1_{ij}{}^{et} f_{\vec{A}}$, $j=0, 5$, $Q_i S^1 \text{et}(A, B)$ transforms A to $S^1_{ij}{}^{et} f_{\vec{A}}$, where A, B may be ordered sets of continual elements, $i=1, 2, 3, 4$.
3. Operations are taking place:

4. ${}^B_D S^1 t_{\vec{A}}^{\vec{A}} = ({}^B_A S^1 t_{\vec{B}}^{\vec{A}})^*$, $\mu({}^B_D S^1 t_{\vec{A}}^{\vec{A}}) = (\mu({}^B_A S^1 t_{\vec{B}}^{\vec{A}}) + 2\mu(B) - \mu(D - A))$
5. ${}^B_A S^1 t_{\vec{B}}^{\vec{A}} = ({}^B_A S^1 t_{\vec{B}}^{\vec{B}})^*$, $\mu({}^B_A S^1 t_{\vec{B}}^{\vec{A}}) = (2\mu(B) - \mu(A) + \mu(A - B))$
6. ${}^B_A S^1 t_{\vec{B}}^{\vec{A}} = ({}^B_A S^1 t_{\vec{B}}^{\vec{A}})^*$, $\mu({}^B_A S^1 t_{\vec{B}}^{\vec{A}}) = (2\mu(B) + \mu(A) - \mu(A - B))$
7. ${}^B_Q S^1 t_{\vec{A}}^{\vec{A}} = ({}^B_Q S^1 t_{\vec{B}}^{\vec{A}})^*$, $\mu({}^B_Q S^1 t_{\vec{A}}^{\vec{A}}) = (2\mu(B) + \mu(A - B) - \mu(Q - B))$

8. ${}^B_S S^1 t_B^A = ({}^B_S S^1 t_B^A)^*$, $\mu({}^B_S S^1 t_B^A) = (2\mu(B) + \mu(A - B) - \mu(Q))$
9. ${}^B_S S^1 t_B^A = ({}^B_S S^1 t_B^A)^*$, $\mu({}^B_S S^1 t_B^A) = (2\mu(B) + \mu(A) - \mu(Q - B))$
10. ${}^B_S S^1 t_B^A = ({}^B_S S^1 t_B^A)^*$, $\mu({}^B_S S^1 t_B^A) = (\mu(B) + \mu(A - B) + \mu(R) - \mu(Q - B))$

Dynamical Continual S¹et-Elements

Also, may be considered dynamical continual S¹et elements, where may be transfer these definitions, operations using on them by analogy [3].

Definition 37

The process of the containment of A(t) into B(t) and the displacement of D(t) from C(t) at time to simultaneously, where some or any elements may be by ordered elements, we shall call dynamical continual S²et – element. Let's denote ${}^{C(t)}_{D(t)} S^1 t(t) {}^{A(t)}_{B(t)}$

Definition 38

The process ${}^{C(t)}_{D(t)} S^1 t(t) {}^{A(t)}_{B(t)}$ is called an ordered dynamical continual S²et – element, if some or any elements from A(t), B(t), C(t), D(t) may be by ordered dynamical continual elements.

It is allowed to multiply dynamical continual S¹et – elements:

$${}^{C_1(t)}_{D_1(t)} S^1 t(t) {}^{A_1(t)}_{B_1(t)} * {}^{C_2(t)}_{D_2(t)} S^1 t(t) {}^{A_2(t)}_{B_2(t)} = {}^{C_1(t)}_{D_1(t) \cup D_2(t)} S^1 t(t) {}^{A_1(t) \cup A_2(t)}_{B_1(t)} (*_4),$$

where some or any elements may be by ordered dynamical continual elements.

Dynamical continual S¹et – elements can be elements of a group by multiplication (*₄).

Dynamical Continual Containment of Oneself

Definition 39

The dynamical continual S¹et -capacity A(t) in itself and from itself of the null type is the process of a containment itself as an element and expelling oneself out of oneself at time to simultaneously, where A(t) – S¹et of dynamical continual elements:

$${}^{A(t)}_{A(t)} S^1 t(t) {}^{A(t)}_{A(t)}. \text{ Denote } S^{1et}_0(t) fA(t).$$

Definition 40

The ordered dynamical continual S¹et -capacity $\overrightarrow{A(t)}$ in itself and from itself of the null type is the capacity containing itself as an element and expelling oneself out of oneself at time t simultaneously, where $\overrightarrow{A(t)}$ – ordered set of dynamical continual elements:

$$\overrightarrow{A(t)} S^1 t(t) \overrightarrow{A(t)}. \text{ Denote } S^{1et}_0(t) f\vec{A}(t)$$

Definition 41

The dynamical continual S¹et -capacity in itself A(t) and from itself B(t) of the first type is the process of a containment of A(t) itself as an element and expelling B(t) oneself out of oneself at time t simultaneously, where A(t), B(t)- sets of dynamical continual elements:

$${}^{B(t)}_{B(t)} S^1 t(t) {}^{A(t)}_{A(t)}. \text{ Denote } S^{1et}_1(t) f_{B(t)}^{A(t)}.$$

Definition 42

The dynamical continual S¹et 1-capacity of the second type is the process of putting B(t) into A(t) and expelling B(t) oneself out of oneself at time t simultaneously, where A(t), B(t)- sets of dynamical continual elements:

$${}^{B(t)}_{B(t)} S^1 t(t) {}^{B(t)}_{A(t)}. \text{ Denote } S^{1et}_2(t) f_{B(t)}^{A(t)}$$

Definition 43

The dynamical continual S¹et 1-capacity of the third type is the process of a containment of B(t) itself as an element and the displacement of B(t) from A(t) at time t simultaneously, where A, B- sets of dynamical continual elements:

$${}^{A(t)}_{B(t)} S^1 t(t) {}^{B(t)}_{B(t)}. \text{ Denote } S^{1et}_3(t) f_{B(t)}^{A(t)}$$

Definition 44

The dynamical continual S¹et -capacity A(t) in itself of the fourth type is the process of a containment of the program that allows it to be generated and it to be degenerated at time t simultaneously, where A(t)- set of dynamical continual elements. Let's denote $S^{1et}_4(t) fA(t)$.

Definition 45

The dynamical continual S¹et -capacity A(t) in itself of the fifth type is the process of a containment of itself in part and expelling oneself in part or contains a program that allows it to be generated and it to be degenerated at time t through the structure S¹et, or both simultaneously, where A(t)- set of dynamical continual elements. Let us denote $S^{1et}_5(t) fA(t)$.

Definition 46

The ordered dynamical continual S¹et -capacity in itself A(t) and from itself $\overrightarrow{B(t)}$ of the first type is the process of a containment of $\overrightarrow{A(t)}$ itself as an element and expelling $\overrightarrow{B(t)}$ oneself out of oneself at time to simultaneously, where A(t)- ordered set of dynamical continual elements, B(t)- set of dynamical continual elements: ${}^{B(t)}_{B(t)} S^1 t(t) {}^{A(t)}_{A(t)}$. Denote $S^{1et}_1(t) f_{B(t)}^{A(t)}$.

Definition 47

The ordered dynamical continual S¹et 1-capacity in itself $\overrightarrow{A(t)}$ and from itself $\overrightarrow{B(t)}$ of the first type is the process of a containment of $\overrightarrow{A(t)}$ itself as an element and expelling $\overrightarrow{B(t)}$ oneself out of oneself at time t simultaneously, where B(t) -ordered set of dynamical continual elements, A- set of dynamical continual elements: $\overrightarrow{B(t)} S^1 t(t) \overrightarrow{A(t)}$. Denote $S^{1et}_1(t) f_{B(t)}^{A(t)}$.

Definition 48

The ordered dynamical continual S¹et 2-capacity in itself $\overrightarrow{A(t)}$ and from itself $\overrightarrow{B(t)}$ of the first type is the process of a containment of $\overrightarrow{A(t)}$ itself as an element and expelling $\overrightarrow{B(t)}$ oneself out of oneself at time t simultaneously, where $\overrightarrow{A(t)}$, $\overrightarrow{B(t)}$ -ordered sets of dynamical continual elements:

$$\overrightarrow{B(t)} S^1 t(t) \overrightarrow{A(t)}. \text{ Denote } S^{1et}_1(t) f_{B(t)}^{A(t)}$$

Definition 49

The dynamical continual S¹et 1-capacity of the second type is the process of a containment B(t) into $\overrightarrow{A(t)}$ and expelling B(t) oneself out of oneself at time to simultaneously, where $\overrightarrow{A(t)}$ - ordered set of dynamical continual elements, B(t) - set of dynamical continual elements:

$$\frac{B(t)}{B(t)} S^1 t(t) \frac{B(t)}{A(t)}. \text{ Denote } S^{1et}_2(t) f_{B(t)}^{\overline{A(t)}}.$$

Definition 50

The dynamical continual S1et 2-capacity of the second type is the process of a containment $\overline{B(t)}$ into $\overline{A(t)}$ and expelling $\overline{B(t)}$ oneself out of oneself at time to simultaneously, where $\overline{B(t)}$ -ordered set of dynamical continual elements, $\overline{A(t)}$ - set of dynamical continual elements:

$$\frac{\overline{B(t)}}{B(t)} S^1 t(t) \frac{\overline{B(t)}}{A(t)}. \text{ Denote } S^{1et}_2(t) f_{B(t)}^{\overline{A(t)}}.$$

Definition 51

The dynamical continual S1et³-capacity of the second type is the process of a containment $\overline{B(t)}$ into $\overline{A(t)}$ and expelling $\overline{B(t)}$ oneself out of oneself at time to simultaneously, where $\overline{A(t)}, \overline{B(t)}$ -ordered sets of dynamical continual elements:

$$\frac{\overline{B(t)}}{B(t)} S^1 t(t) \frac{\overline{B(t)}}{A(t)}. \text{ Denote } S^{1et}_2(t) f_{B(t)}^{\overline{A(t)}}.$$

Definition 52

The dynamical continual S1et¹-capacity of the third type is the process of a containment of $\overline{B(t)}$ itself as an element and the displacement of $B(t)$ from $\overline{A(t)}$ at time t simultaneously, where $\overline{A(t)}$ ordered set of dynamical continual elements, $B(t)$ - set of dynamical continual elements:

$$\frac{\overline{A(t)}}{B(t)} S^1 t(t) \frac{B(t)}{B(t)}. \text{ Denote } S^{1et}_3(t) f_{B(t)}^{\overline{A(t)}}.$$

Definition 53

The dynamical continual S1et 2-capacity of the third type is the process of a containment of $\overline{B(t)}$ itself as an element and the displacement of $B(t)$ from $\overline{A(t)}$ at time t simultaneously, where $\overline{B(t)}$ ordered set of dynamical continual elements, $\overline{A(t)}$ - set of dynamical continual elements:

$$\frac{\overline{A(t)}}{B(t)} S^1 t(t) \frac{\overline{B(t)}}{B(t)}. \text{ Denote } S^{1et}_3(t) f_{B(t)}^{\overline{A(t)}}.$$

Definition 54

The dynamical continual S1et 3-capacity of the third type is the process of a containment of $\overline{B(t)}$ itself as an element and the displacement of $\overline{B(t)}$ from $\overline{A(t)}$ at time t simultaneously, where $\overline{A(t)}, \overline{B(t)}$ -ordered sets of dynamical continual elements:

$$\frac{\overline{A(t)}}{B(t)} S^1 t(t) \frac{\overline{B(t)}}{B(t)}. \text{ Denote } S^{1et}_3(t) f_{B(t)}^{\overline{A(t)}}.$$

Definition 55

The ordered dynamical continual S1et -capacity $\overline{A(t)}$ in itself of the fourth type is the process that contains the program that allows it to be generated and it to be degenerated at time to simultaneously, where $\overline{A(t)}$ - S1et of dynamical continual elements. Let's denote

$$S^{1et}_4(t) f_{\overline{A(t)}}$$

Definition 56

The ordered dynamical continual S1et -capacity $\overline{A(t)}$ in itself of the fifth type is the process of a containment of itself in part and expelling oneself in part or contains a program that allows it to be generated in part and it to be degenerated in part at time

t , or both simultaneously, where $\overline{A(t)}$ - ordered S1et of dynamical continual elements. Let us denote

$$S^{1et}_5(t) f_{\overline{A(t)}}$$

Also we consider some elements:

$$S^{1et}_0(t) f_{\uparrow I \downarrow -1}, S^{1et}_1(t) f_{B(t) \uparrow \downarrow -1}, S^{1et}_2(t) f_{B(t) \uparrow \downarrow -1},$$

$$S^{1et}_3(t) f_{B(t) \uparrow \downarrow -1}, S^{1et}_4(t) f_{\uparrow I \downarrow -1}$$

[3] and etc.

Dynamical Continual Mathematics S1et-Itself

- Similarly, for the simultaneous execution of various operators:

$$F_0(t)C(t)S^1 t(t)F_2(t)A(t) \\ F_1(t)D(t)F_3(t)B(t)$$

are operators.

- Similarly, for the simultaneous execution of various operators:

$$S^{1et}_j(t) f_{F(t)A(t)} \quad j=0,4,5, \text{ and } S^{1et}_k(t) f_{B(t)}^{A(t)} \quad k=1,2,3, \text{ where } \{F(t)\} =$$

$$(F_0(t), F_1(t), F_2(t), F_3(t))$$

are operators.

$$3. \quad \frac{B(t)}{D(t)} S^1 t(t) \frac{A(t)}{B(t)} = \left(\frac{B(t)}{A(t)} S^1 t(t) \frac{A(t)}{B(t)} * \right) \mu_{D(t)-A(t)}^{B(t)} S^1 t(t) \frac{A(t)}{B(t)} =$$

$$\left(\mu^{ss} \left(\frac{B(t)}{A(t)} S^1 t(t) \frac{A(t)}{B(t)} * \right) \right) \\ \mu(A(t)) + 2\mu(B(t)) - \mu(D(t) - A(t))$$

$$4. \quad \frac{B(t)}{A(t)} S^1 t(t) \frac{A(t)}{B(t)} = \left(\frac{B(t)}{A(t)} S^1 t(t) \frac{B(t)}{B(t)} * \right) \\ \frac{B(t)}{A(t)} S^1 t(t) \frac{A(t)-B(t)}{B(t)},$$

$$5. \quad \mu_{A(t)}^{B(t)} S^1 t(t) \frac{A(t)}{B(t)} = \left(\mu^{ss} \left(\frac{B(t)}{A(t)} S^1 t(t) \frac{B(t)}{B(t)} * \right) \right) \\ 2\mu(B(t)) + \mu(A(t) - B(t)) - \mu(A(t))$$

$$6. \quad \frac{B(t)}{A(t)} S^1 t(t) \frac{A(t)}{B(t)} = \left(\frac{B(t)}{B(t)} S^1 t(t) \frac{A(t)}{B(t)} * \right) \\ \frac{B(t)}{A(t)-B(t)} S^1 t(t) \frac{A(t)}{B(t)},$$

$$7. \quad \mu_{A(t)}^{B(t)} S^1 t(t) \frac{A(t)}{B(t)} = \left(\mu^{ss} \left(\frac{B(t)}{B(t)} S^1 t(t) \frac{A(t)}{B(t)} * \right) \right) \\ 2\mu(B(t)) + \mu(A(t)) - \mu(A(t) - B(t))$$

$$8. \quad \frac{B(t)}{Q(t)} S^1 t(t) \frac{A(t)}{B(t)} = \left(\frac{S^{1et}_0 f(t) B(t)}{Q(t)-B(t)} S^1 t(t) \frac{A(t)-B(t)}{B(t)} * \right) \\ \mu_{Q(t)-B(t)}^{B(t)} S^1 t(t) \frac{A(t)}{B(t)} =$$

$$\left(\mu^{ss} \left(\frac{S^{1et}_0 f(t) B(t)}{Q(t)-B(t)} S^1 t(t) \frac{A(t)}{B(t)} * \right) \right) \\ 2\mu(B(t)) + \mu(A(t) - B(t)) - \mu(Q(t) - B(t))$$

$$9. \quad \frac{B(t)}{Q(t)} S^1 t(t) \frac{A(t)}{B(t)} = \left(\frac{B(t)}{Q(t)} S^1 t(t) \frac{B(t)}{B(t)} * \right) \\ \frac{B(t)}{Q(t)} S^1 t(t) \frac{A(t)-B(t)}{B(t)},$$

$$\mu_{Q(t)}^{B(t)} S^1 t(t) \frac{A(t)}{B(t)} = \left(\mu^{ss} \left(\frac{B(t)}{Q(t)} S^1 t(t) \frac{B(t)}{B(t)} * \right) \right) \\ 2\mu(B(t)) + \mu(A(t) - B(t)) - \mu(Q(t))$$

$$10. \quad \frac{B(t)}{Q(t)} S^1 t(t) \frac{A(t)}{B(t)} = \left(\frac{B(t)}{B(t)} S^1 t(t) \frac{A(t)}{B(t)} * \right) \\ \frac{B(t)}{Q(t)-B(t)} S^1 t(t) \frac{A(t)}{B(t)},$$

$$\mu_{Q(t)}^{B(t)} S^1 t(t) \frac{A(t)}{B(t)} = \left(\mu^{ss} \left(\frac{B(t)}{B(t)} S^1 t(t) \frac{A(t)}{B(t)} * \right) \right) \\ 2\mu(B(t)) + \mu(A(t)) - \mu(Q(t) - B(t))$$

$$11. \quad \begin{aligned} \frac{R(t)}{Q(t)} S^1 t(t)_{B(t)}^{A(t)} &= \left(\begin{array}{c} S_3^{1et} f(t)_{R(t)}^{B(t)} * \\ R(t) S^1 t(t)_{B(t)}^{A(t)-B(t)} \end{array} \right) \\ \mu_{Q(t)}^{(R(t))} S^1 t(t)_{B(t)}^{A(t)} &= \left(\begin{array}{c} \mu^{ss} (S_3^{1et} f(t)_{R(t)}^{B(t)} *) \\ \mu(B(t)) + \mu(A(t) - B(t)) + \mu(R(t)) - \mu(Q(t) - B(t)) \end{array} \right) \end{aligned}$$

Connection of Dynamical Continual S²et-Elements with Dynamical Containment of Oneself

Consider a fifth type of dynamical partial containment of oneself. For example, $S_5^{1et}(t)f_{A^n(t)}^{A^n(t)}$, where $\{A^n(t)\} = \{a_1(t), a_2(t), \dots, a_n(t)\}$, i.e. n - continual elements, it is possible to consider the dynamical containment of oneself $S_5^{1et}(t)f_{A^m(t)}^{A^m(t)}$ with m continual elements from $\{A^n(t)\}$, at $m < n$, which is process to be formed by the form (1) [1], that is, only m continual elements from $\{A^n(t)\}$ are located in the structure $S_5^{1et}(t)f_{A^m(t)}^{A^m(t)}$. Dynamical continual containments of oneself of the fifth type can be formed for any other structure, not necessarily S1et, only through the obligatory reduction in the number of continual elements in the structure. In particular, using the form (2) [2]. Structures more complex than $S_5^{1et}(t)f_{A^n(t)}^{A^n(t)}$, can be introduced.

Dynamical Continual S¹et-Elements with Target Weights

Also, may be considered dynamical continual S1et -elements with target weights, where may be transfer these definitions, operations using on them by analogy [3].

Definition 57

The process of the containment of $A(t)$ with target weights $\{g_1(t)\}$ into $B(t)$ and the displacement of $D(t)$ with target weights $\{g_2(t)\}$ from $C(t)$ at time t simultaneously, where some or any elements may be by dynamical continual elements, we shall call dynamical continual S2et – element with target weights. Let's denote $\frac{C(t)}{D(t)\{g_2(t)\}} S^1 t(t)_{B(t)}^{A(t)\{g_1(t)\}}$.

Definition 58

The process $\frac{C(t)}{D(t)\{g_2(t)\}} S^1 t(t)_{B(t)}^{A(t)\{g_1(t)\}}$ is called an ordered dynamical continual S1et – element with target weights $\{g_1(t)\}$ or $\{g_2(t)\}$ at time t , or both simultaneously, if some or any elements from A, B, C, D may be by ordered dynamical continual elements with target weights.

It is allowed to multiply dynamical continual S1et –elements with target weights:

$$\begin{aligned} \frac{C_1(t)}{D_1(t)\{g_2(t)\}} S^1 t(t)_{B_1(t)}^{A_1(t)\{g_1(t)\}} * \frac{C_2(t)}{D_2(t)\{g_2(t)\}} S^1 t(t)_{B_2(t)}^{A_2(t)\{g_1(t)\}} = \\ \frac{C_1(t)}{(D_1(t) \cup D_2(t))\{g_2(t)\}} S^1 t(t)_{B_1(t)}^{(A_1(t) \cup A_2(t))\{g_1(t)\}} (*_5), \end{aligned}$$

where some or any elements may be by ordered dynamical continual elements with target weights or dynamical continual elements with target weights, or both.

Dynamical continual S1et – elements with target weights can be elements of a group by multiplication (*5).

Dynamical Continual Containment of Oneself with Target Weights

Definition 59

The dynamical continual S1et -capacity $A(t)$ in itself and from itself with target weights $\{g(t)\}$ of the null type is the process

of a containment itself as an element with target weights $\{g(t)\}$ and expelling oneself out of oneself with target weights $\{g(t)\}$ at time t simultaneously, where $A(t)$ – S1et of some dynamical continual elements or some ordered dynamical continual elements, or both. Denote $S_0^{1et}(t)f_{A(t)}\{g(t)\}$

Definition 60

The dynamical continual S2et -capacity $A(t)$ in itself with target weights $\{g(t)\}$ of the fourth type is the process that contains the program that allows it to be generated with target weights $\{g(t)\}$ and it to be degenerated with target weights $\{g(t)\}$ at time t simultaneously, where $A(t)$ – S1et of some dynamical continual elements or some ordered dynamical continual elements, or both. Denote $S_4^{1et}(t)f_{A(t)}\{g(t)\}$.

Definition 61

The ordered dynamical continual S1et -capacity $A(t)$ in itself of the fifth type with target weights $\{g(t)\}$ is the process of a containment of itself in part with target weights $\{g(t)\}$ and expelling oneself in part with target weights $\{g(t)\}$ or contains a program that allows it to be generated in part with target weights $\{g(t)\}$ and it to be degenerated in part with target weights $\{g(t)\}$ at time t simultaneously, or both simultaneously, where $A(t)$ - Set of some dynamical continual elements or some ordered dynamical continual elements, or both. Denote

$$S_5^{1et}(t)f_{A(t)}\{g(t)\}$$

Definition 62

The dynamical continual S1et -capacity in itself with target weights $\{g_1(t)\}$ and from itself $B(t)$ with target weights $\{g_2(t)\}$ of the first type is the process of a containment of $A(t)$ itself as an element $A(t)$ and expelling $B(t)$ oneself with target weights $\{g_2(t)\}$ out of oneself at time t simultaneously, where some or any elements from $A(t), B(t)$ may be by ordered dynamical continual elements with target weights or dynamical continual elements with target weights, or both:

$$\frac{B(t)}{t\{g_2(t)\}} S^1 t(t)_{A(t)}^{A(t)\{g_1(t)\}}. \text{ Denote } S_1^{1et}(t)f_{B(t)\{g_2(t)\}}^{A(t)\{g_1(t)\}}.$$

Definition 63

The dynamical continual S1et 1-capacity with target weights of the second type is the process of putting $B(t)$ with target weights $\{g_1(t)\}$ into $A(t)$ and expelling $B(t)$ oneself with target weights $\{g_2(t)\}$ out of oneself at time t simultaneously, where some or any elements from $A(t), B(t)$ may be by ordered dynamical continual elements with target weights or dynamical continual elements with target weights, or both:

$$\frac{B(t)}{B(t)\{g_2(t)\}} S^1 t(t)_{A(t)}^{B(t)\{g_1(t)\}}. \text{ Denote } S_2^{1et}(t)f_{B(t)\{g_1(t)\},\{g_2(t)\}}^{A(t)}.$$

Definition 64

The dynamical continual S1et 1-capacity of the third type with target weights is the process of a containment of $B(t)$ itself as an element with target weights $\{g_1(t)\}$ and the displacement of $B(t)$ with target weights $\{g_1(t)\}$ from $A(t)$ at time t simultaneously, where some or any elements from $A(t), B(t)$ may be by ordered dynamical continual elements with target weights or dynamical continual elements with target weights, or both:

$$\frac{A(t)}{B(t)\{g_1(t)\}} S^1 t(t)_{B(t)}^{B(t)\{g_1(t)\}}. \text{ Denote } S_3^{1et}(t)f_{B(t)\{g_1(t)\}}^{A(t)}$$

Also we consider some elements:

$$S_0^{1et}(t) f \uparrow \downarrow \uparrow \downarrow \{g(t)\}, S_1^{1et}(t) f_{B(t) \uparrow \downarrow \infty}^{11 \uparrow \downarrow \{g(t)\}}, S_2^{1et}(t) f_{B(t) \uparrow \downarrow \infty}^{A(t) \uparrow \downarrow \{g(t)\}}, S_3^{1et}(t) f_{B(t) \uparrow \downarrow \infty}^{A(t) \uparrow \downarrow \{g(t)\}}, S_4^{1et}(t) f \uparrow \downarrow \uparrow \downarrow \{g(t)\} \text{ etc.}$$

Mathematics Itself for Dynamical Continual S'et-Elements with Target Weights

1. Similarly, for the simultaneous execution of various operators:

$$F_0(t)C(t)S_1^{1et}(t)F_2(t)A(t) \\ F_1(t)D(t)g_2(t)F_3(t)B(t)$$

where $F_0(t), F_1(t), F_2(t), F_3(t)$ are operators.

2. Similarly, for the simultaneous execution of various operators:

$$S_j^{1et}(t) f F(t) A(t), j=0,4,5, \text{ and } S_k^{1et}(t) f_{B(t)g_2(t)}^{A(t)},$$

$k=1,2,3$, where $\{F(t)\} = (F_0(t), F_1(t), F_2(t), F_3(t))$ are operators.

$$\begin{aligned} 1. \quad & \frac{B(t)}{D(t)g_1(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) = \left(\frac{B(t)}{A(t)g_1(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) * \right. \\ & \left. \frac{B(t)}{(D(t)-A(t))g_1(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) \right) \cdot \mu \left(\frac{B(t)}{D(t)g_1(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) \right) = \\ & \left(\mu^s \left(\frac{B(t)}{A(t)g_1(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) * \right) \right) \\ & \left(\mu(A(t)g_1(t)) + 2\mu(B(t)) - \mu((D(t)-A(t))g_1(t)) \right) \\ 2. \quad & \frac{B(t)}{A(t)g_1(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) = \left(\frac{B(t)}{A(t)g_1(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) * \right. \\ & \left. \frac{B(t)}{A(t)g_1(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) \right) \cdot \mu \left(\frac{B(t)}{A(t)g_1(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) \right) = \\ & \left(\mu^{ss} \left(\frac{B(t)}{A(t)g_1(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) * \right) \right) \\ & \left(2\mu(B(t)) + \mu(A(t)-B(t)) - \mu(A(t)g_1(t)) \right) \\ 3. \quad & \frac{B(t)}{A(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) = \left(\frac{B(t)}{A(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) * \right. \\ & \left. \frac{B(t)}{A(t)-B(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) \right) \cdot \mu \left(\frac{B(t)}{A(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) \right) = \\ & \left(\mu^{ss} \left(\frac{B(t)}{A(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) * \right) \right) \\ & \left(2\mu(B(t)) + \mu(A(t)g_1(t)) - \mu(A(t)-B(t)) \right) \\ 4. \quad & \frac{B(t)}{Q(t)g_1(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) = \left(\frac{S_0^{1et}(t) f(t) B(t) g_1(t) *}{(Q(t)-B(t))g_1(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) \right) \cdot \\ & \mu \left(\frac{B(t)}{Q(t)g_1(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) \right) = \left(\mu^{ss} \left(S_0^{1et}(t) f(t) B(t) g_1(t) * \right) \right) \\ & \left(2\mu(B(t)) + \mu((A(t)-B(t))g_1(t)) - \mu((Q(t)-B(t))g_1(t)) \right) \\ 5. \quad & \frac{B(t)}{Q(t)g_2(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) = \left(\frac{B(t)}{Q(t)g_2(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) * \right. \\ & \left. \frac{B(t)}{Q(t)g_2(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) \right) \cdot \mu \left(\frac{B(t)}{Q(t)g_2(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) \right) = \\ & \left(\mu^{ss} \left(\frac{B(t)}{Q(t)g_2(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) * \right) \right) \\ & \left(2\mu(B(t)) + \mu((A(t)-B(t))g_1(t)) - \mu(Q(t)g_2(t)) \right) \\ 6. \quad & \frac{B(t)}{Q(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) = \left(\frac{B(t)}{Q(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) * \right. \\ & \left. \frac{B(t)}{Q(t)-B(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) \right) \cdot \mu \left(\frac{B(t)}{Q(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) \right) = \\ & \left(\mu^{ss} \left(\frac{B(t)}{Q(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) * \right) \right) \\ & \left(2\mu(B(t)) + \mu(A(t)g_1(t)) - \mu(Q(t)-B(t)) \right) \\ 7. \quad & \frac{R(t)g_1(t)}{Q(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) = \left(\frac{S_3^{1et}(t) f(t) R(t)g_1(t) *}{(Q(t)-B(t))g_1(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) \right) \cdot \\ & \mu \left(\frac{R(t)g_1(t)}{Q(t)} S_1^{1et}(t) \frac{A(t)}{B(t)} g_1(t) \right) = \left(\mu^{ss} \left(S_3^{1et}(t) f(t) R(t)g_1(t) * \right) \right) \\ & \left(\mu(B(t)) + \mu(A(t)-B(t)) + \mu(R(t)g_1(t)) - \mu(Q(t)-B(t)) \right) \end{aligned}$$

Connection of Dynamical Continual S'et-Elements with Target Weights with Dynamical Containment of Oneself with Target Weights

Consider a fifth type of dynamical partial containment of oneself with target weights $g(t)$. For example, based on

$$S_5^{1et}(t) f A(t) \{g(t)\},$$

where $A=(a_1(t), a_2(t), \dots, a_n(t))$, i.e. n - continual elements with target weights $\{g(t)\}$ in one point x , it is possible to consider the dynamical containment of oneself with target weights

$$S_5^{1et}(t) f A(t) \{g(t)\},$$

with m continual elements with target weights $\{g(t)\}$ from A , at $m < n$, which is process to be formed by the form (1), that is, only m continual elements with target weights $\{g(t)\}$ from A are located in the structure $S_5^{1et}(t) f A(t) \{g(t)\}$,

Dynamical containments of oneself with target weights of the fifth type can be formed for any other structure, not necessarily S2et, only through the obligatory reduction in the number of continual elements with target weights in the structure. In particular, using the form (2). Structures more complex than

$$S_5^{1et}(t) f A(t) \{g(t)\},$$

can be introduced.

Supplement

We consider S'et-logic: consider the functional $f(Q)$, which gives a numerical value for the truth of the statement Q from the interval $[0,1]$, where 0 corresponds to "no", and 1 corresponds to the logical value "yes". Then for joint statements

$$\begin{aligned} & f(\frac{C}{D} S_1^A t) + f(S_1^A t_B) - f(\frac{C}{D} S_1^A t \cap S_1^A t_B) = \\ & \left(f^{os1}(C \cap D - Co(C \cap D)) + f^{s1}(A \cap B) + f^{-s1}(C - Co(D) S_1^A t) \right) \cdot f(\frac{C}{D} S_1^A t \cap S_1^A t_B), f^{s1}(x) \\ & f(A) + f(B) - f(A \cap B) + f(D) - f(C) \end{aligned}$$

the value of self-truth for self-statement x , $Co(x)$ - content of x , $f^{os1}(x)$ - the value of oself-truth for oself-statement x ; for dependent statements: $f(A*B)=f(A)*f(B/A)=f(B)*f(A/B)$, where $f(B/A)$ - conditional truth of the statement B at the statement A , $f(A/B)$ - conditional truth of the statement A at the statement B , for dependent statements:

$$f(\frac{C}{D} S_1^A t \cap S_1^A t_B) = f(\frac{C}{D} S_1^A t) * f(S_1^A t_B / \frac{C}{D} S_1^A t) = f(S_1^A t_B) * f(\frac{C}{D} S_1^A t / S_1^A t_B).$$

Adding the truth values of inconsistent propositions: $f(A+B)=F(A)+f(B)$. The formula of complete truth:

$$f(A) = \sum_{k=1}^n f(B_k) * f(A/B_k),$$

B_1, B_2, \dots, B_n -full group of hypotheses-statements:

$$\sum_{k=1}^n f(B_k) = 1 \text{ ("yes")}. \quad \square$$

Remark

A statement can be interpreted as an event, and its truth value as a probability.

Definition 65. S'et-probability of events

$$\frac{C}{D} S_1^A t_B \text{ is } p(\frac{C}{D} S_1^A t_B), \text{ denote } \frac{C}{D} S_1^A p_B^A.$$

In particular, $\frac{C}{D} S_1^A p_B^A$ for joint events $S_1^A t_B, \frac{C}{D} S_1^A t, A, B, C, D, \frac{C}{D} S_1^A t_B$:

$$p(\frac{C}{D} S_1^A t_B) = p(\frac{C}{D} S_1^A t) + p(S_1^A t_B) - p(S_1^A t_B \cap \frac{C}{D} S_1^A t) =$$

$$\left(p^{os1}(C \cap D - Co(C \cap D)) + p^{s1}(A \cap B) + p^{-s1}(C - Co(D) S_1^A t) \right) \cdot p(S_1^A t_B \cap \frac{C}{D} S_1^A t), \\ p(A) + p(B) - p(A \cap B) + p(D) - p(C)$$

for dependent events:

$$p(\frac{C}{D}S^1t \cap S^1t_B^A) = p(\frac{C}{D}S^1t) * p(S^1t_B^A / \frac{C}{D}S^1t) = p(S^1t_B^A) * p(\frac{C}{D}S^1t / S^1t_B^A).$$

$p^{s1}(x)$ - the value of self-P for self- event x , $Co(x)$ – content of x , $p^{os1}(x)$ - the value of oself-P for oself- event x .

Energy of a Living Organism

$$f(r, a(E_q)) = St_{t_0}^{\{q(\frac{a}{a}St_a^a) St_q^{Eq} W_q(\frac{a}{a}St_a^a)\}, St_{dr}^{\{El^{dr}, q(\frac{a}{a}St_a^a)\}}}$$

$\frac{a}{a}St_a^a$ internal energy of a living organism, q - a gap in the energy cocoon of a living organism, r -the position of the assemblage point dr on the energy cocoon of a living organism, W_q - energy prominences from the gap in the cocoon of a living organism, Eq -external energy entering the gap in the cocoon of a living organism, El^{dr} - a bundle of fibers of external energy self-capacities, collected at the point of assembly of the cocoon of a living organism.

$\frac{A}{A}St$ will be called anti-capacity from oneself. For example, “white hole” in physics is such simple anti-capacity. The concepts of “white hole” and “black hole” were formulated by the physicists proceeding from the physics subjects –usual energies level. The mathematics allows to find deeply and to formulate the concepts singular points in the Universe proceeding from levels of more thin energies. The experiments of Nobel laureates in 2022-year Asle Ahlen, Clauser John, Zeilinger Anton correspond to the concept of the Universe as its self-containment in itself. The connection between the elements of self-containment in itself is a property of self-containment in itself and therefore does not disappear when their location in it changes. The energy of self-containment in itself is closed on itself.

Hypothesis

The containment of the galaxy in oneself as spiral curl and the expelling her out of oneself defines its existence. A self-consistency in itself as an element A is the god of A , the self-consistency in itself as an element the globe—the god of the globe, the self-consistency in itself as an element man-- the god of the man, the self-consistency in itself as an element of the universe-- the god of the universe, the containment of A into oneself is spirit of A , the containment of the globe into oneself is spirit of globe, the containment of the man into oneself is spirit of the man (soul), the containment of the universe into oneself is spirit of the universe. We may consider the next axiom: any holding capacity is the capacity of oneself in itself. This is for each energy capacity. The Chinese book of Changes “I Ching” uses a structure similar to (*) implicitly.

Using elements of the mathematics of Sit [2]. we introduce the concept of Sit – the change in physical quantity B : $St_x^{\{B_1, B_2, \dots, B_n\}}$. Then the mean Sit - velocity will be $v_{sit}(t, \Delta t) = St_x^{\{B_1, B_2, \dots, B_n\}} / \Delta t$, and Sit-velocity at time $t: v_{sit} = \lim_{\Delta t \rightarrow 0} v_{qst}(t, \Delta t) Sit$ – acceleration $a_{st} = \frac{dv_{sit}}{dt}$.

In normal use, simply Sitx reduce to result a sum at point x of space, and when using Sitx with "target weights", we get, depending on the "target weights", one or another modification, namely, for example, the velocity v_{st}^f (with a "target weight" f)

in the case when two velocities v_1, v_2 are involved in the set $\{v_1, f, v_2\}$ for $v_{st}^f = St_x^{\{v_1, f, v_2\}}$, f – instantaneous replacement we get an instantaneous substitution v_1 by v_2 at point x of space at time t_0 .

Consider, in particular, some examples: 1) $St_{\{x, 1, x, 2\}}^e$ describes the presence of the same electron e at two different points x_1, x_2 . 2) The nuclei of atoms can be considered as Sit elements.

Similarly, the concepts of Sit - force, Sit - energy are introduced. For example, $E_{st}^f = St_x^{\{E1, f, E2\}}$ it would mean the instantaneous replacement of energy $E1$ by $E2$ at time t_0 . Two aspects of Sit– energy should be distinguished: 1) carrying out the desired "target weight", 2) the fixing result of it. Do not confuse energy - Sit (this is the node of energies) with Sit – energy that generates the node of energies, usually with the "target weights". In the case of ordinary energies, the energy node is carried out automatically. In fact, sit – elements are all ordinary, but with "target weights" they become peculiar. Here you need the necessary kind of energy to perform them. As a rule, this energy lies in the region itself. This is natural, since it's much easier to control the elements of the k level by the elements of the more highly structured $k + 1$ level. Consider the concepts of capacity in itself of physical objects. The question arises about the self-energy of the object. In particular, according to the results of the publication [4]. « St_B^B will mean $S_f f B$.» In particular, it allows you to determine the self-energy of DNA through St_{DNA}^{DNA}, St_Q^Q - self-energy Q . The law of self-energy conservation acts on the level of self-energy already.

Supplement for Quantum Mechanics and through Sit-elements: Self-equation Schrödinger type

$$St_{\frac{\partial \hat{\rho}}{\partial t} + [\hat{W}, \hat{\rho}] = 0}, \hat{\rho} = \exp(i\hat{H}_0 t / \hbar) \hat{\rho} \exp(-i\hat{H}_0 t / \hbar), \hat{W} = \exp(i\hat{H}_0 t / \hbar) \hat{W} \exp(-i\hat{H}_0 t / \hbar).$$

Hamilton operator $\hat{H} = \hat{H}_0 + \hat{W}_0$, \hat{H}_0 -considered quantum system energy, consisting of two or more parts, without their interaction with each other, \hat{W}_0 is the energy of their interaction, ρ -statistical operator [3]. Self-energy

$$St_{\hat{H}}^{\hat{H}} = St_{\hat{H}_0 + \hat{W}_0}^{\hat{H}_0 + \hat{W}_0} = St_{\hat{H}_0 + \hat{W}_0}^{\hat{H}_0} + St_{\hat{H}_0 + \hat{W}_0}^{\hat{W}_0} \\ = St_{\hat{H}_0}^{\hat{H}_0} + St_{\hat{W}_0}^{\hat{H}_0} + St_{\hat{H}_0}^{\hat{W}_0} + St_{\hat{W}_0}^{\hat{W}_0}, St_{\hat{H}_0}^{\hat{H}_0}$$

considered quantum system self-energy $St_{\hat{W}_0}^{\hat{W}_0}$ is self-energy of their interaction, $St_{\hat{W}_0}^{\hat{W}_0}$ -object manifestation of the energy of the system in an external field, $St_{\hat{W}_0}^{\hat{H}_0}$ the manifestation of the energy of the system in the energy interaction with the external field. Variants of the Schrödinger equation $\frac{\partial \hat{\rho}}{\partial t} + [\hat{W}, \hat{\rho}] = 0$ of the form S2f, S3f are possible, using the form (1) or form (2) [1]. For Classical statistical Mechanics self-analogue of the equation

$$\frac{\partial \rho}{\partial t} + [H, \rho] = 0 (*) : St_{\frac{\partial \rho}{\partial t} + [H, \rho] = 0}^{\frac{\partial \rho}{\partial t} + [H, \rho] = 0}$$

The carrier of the measure of objectivity-mass should be objectivity-elementary particle graviton, i.e. have the form

$$St_{objectivity}^{objectivity},$$

therefore it is a self-particle and is not an element of the level of objectivity, but is an element of the level self. Therefore,

it cannot be found at our level. In fact, the theory of Sit-elements helps to form a unified field theory on a qualitative level, because it is not possible to create a quantitative unified field theory. Supplement for string theory: May be to try represent elementary particles in the form of continual self-elements of the type

$S_{\infty}^{-} = \sin(-\infty) \rightarrow \downarrow \uparrow \uparrow_{-1}^1, T_{\infty}^{+} = \text{tg} \infty \rightarrow \uparrow \downarrow \downarrow_{\infty}^{\infty}, T_{\infty}^{-} = \text{tg}(-\infty) \rightarrow \downarrow \uparrow \uparrow_{\infty}^{\infty}, f \uparrow \downarrow g$ for any f, g etc.

Similarly, you can consider all this through S^1 et, only in much more interesting versions. But this is already in the next articles. The transition process in the form of $\mathbb{C}St_B^A$ is switched on during the transition from one world A (spatial variables, which we denote by X1, and temporal variables, through T1) to another B (spatial variables, which we denote by X2, and temporal variables, through T2). It is accompanied by spatial variables in form (T1, X1), and temporary - T3.

Declarations

Availability of Data and Material

1. Danilishyn I.V. Danilishyn O.V. THE USAGE OF SIT-ELEMENTS FOR NETWORKS. IV International Scientific and Practical Conference "GRUNDLAGEN DER MODERNEN WISSENSCHAFTLICHEN FORSCHUNG", 31.03.2023/Zurich, Switzerland. <https://archive.logos-science.com/index.php/conference-proceedings/issue/view/9>
2. Danilishyn I.V. Danilishyn O.V. MATHEMATICS ST, PROGRAMMING OPERATORS ST AND SOME EMPLOYMENT. Collection of scientific papers "SCIENTIA", 2023. <https://previous.scientia.report/index.php/archive/issue/view/07.04.2023>
3. Dnilishyn I.V. Danilishyn O.V. DYNAMICAL SIT-ELEMENTS . IV International Scientific and Practical Conference "GRUNDLAGEN DER MODERNEN WISSENSCHAFTLICHEN FORSCHUNG ", 31.03.2023/Zurich, Switzerland. <https://archive.logos-science.com/index.php/conference-proceedings/issue/view/9>

Competing Interest

There are no competing interests. All sections of the article are executed jointly.

Funding

There were no sources of funding for writing the article.

Authors' Contributions

The contribution of the authors is the same, we will not separate.

Acknowledgements

We are grateful to our academic supervisor Scientific director: Volodymyr Pasynkov PhD of physic-mathematical science, assistant professor of applied mathematics and calculated techniques department of "National Metallurgical Academy" (Ukraine) for guidance and assistance in writing this article.

References

1. Thomas Jech J (1971) LECTURES IN SET THEORY WITH PARTICULAR EMPHASIS ON THE METHOD OF FORCING. SPRINGER-VERLAG Berlin. Heidelberg.
2. Danilishyn IV, Danilishyn OV (2023) THE USAGE OF SIT-ELEMENTS FOR NETWORKS. IV International Scientific and Practical Conference GRUNDLAGEN DER MODERNEN WISSENSCHAFTLICHEN FORSCHUNG, 31.03.2023/Zurich, Switzerland 129-134.
3. Danilishyn IV, Danilishyn OV (2023) DYNAMICAL SIT-ELEMENTS. IV International Scientific and Practical Conference GRUNDLAGEN DER MODERNEN WISSENSCHAFTLICHEN FORSCHUNG, Zurich, Switzerland. <https://archive.logos-science.com/index.php/conference-proceedings/issue/view/9>
4. Danilishyn IV, Danilishyn OV (2023) MATHEMATICS ST, PROGRAMMING OPERATORS ST AND SOME EMPLOYMENT. Collection of scientific papers "SCIENTIA", <https://previous.scientia.report/index.php/archive/issue/view/07.04.2023>