

An Insight into Tungiasis: Prevalence, Risk Factors And Prevention

Valeri Velev^{1,*}, Viktoria Vicheva¹, Toma Tomov², Yordanka Mitova-Mineva¹

¹Department of Epidemiology and Hygiene, Faculty of Medicine, Medical University of Sofia, Bulgaria

²University Hospital "Prof. Iv. Kirov", Medical University of Sofia, Bulgaria

*Corresponding author: Valeri Velev, Department of Epidemiology and Hygiene, Faculty of Medicine, Medical University of Sofia, Bulgaria.

Submitted: 11 August 2025 Accepted: 18 August 2025 Published: 25 August 2025

doi <https://doi.org/10.63620/MKWJCD.2025>.

Citation: Velev, V., Vicheva, V., Tomov, T., & Mitova-Mineva, Y. (2025). *An Insight into Tungiasis: Prevalence, Risk Factors And Prevention. Wor Jour of Clin Der*, 2(4), 01-05.

Abstract

Introduction: Tungiasis is a neglected parasitic skin disease caused by the permanent penetration of the female sand flea (*Tunga penetrans*) in the skin of the host, mainly the lower extremities. This gradually, it can lead to severe limb deformities. Main endemic areas are sub-Saharan Africa and South America. The purpose is to provide a better understanding of the spread and risk factors leading to infection among locals and tourists in these areas.

Materials and Methods: We conducted a review of the available scientific literature on PubMed, Google Scholar, AJOL, Google Scholar, Web of Science which included randomized controlled trials, meta-analysis, observational studies, prospective controlled studies and case reports.

Results and Discussion: We have established the prevalence of tungiasis and its associated risk factors for the main endemic areas of the world. The risks are associated with age, less often gender; ground floor; infrequent wearing of shoes, contact with animals and residence in rural areas. They are also relevant for tourists in these areas, who can transmit the disease to new communities. It is curious how quickly the disease can spread among people in an environment with limited resources and poor hygiene.

Conclusion: This review shows that there are still large knowledge gaps regarding the prevalence and magnitude of tungiasis in South America and sub-Saharan Africa. It is important to identify the risk factors and the main animal species acting as reservoirs.

Keywords: Tungiasis, Tunga Penetrans, Neglected Tropical Diseases, Epidemiology

Introduction

Tungiasis is one of the neglected tropical parasitic skin diseases. It is caused by the invasion of the female sand flea *Tunga penetrans* into the skin, mainly on the soles of the lower limbs. Very rarely the infestation is caused by a related flea, *Tunga trimamillata*. To date it is a health issue mainly in sub-Saharan Africa, the Caribbean, Latin and South America. It occurs mainly in poor rural populations and is also a major problem for children. In

diurnal zones, the incidence can reach over 73%. Early removal of the parasite from the skin is still the most common therapy. Tungiasis is one of the neglected tropical diseases (DNDi) already included in the World Health Organization's (WHO) DNDi Roadmap 2021-2030, and accordingly surveillance of its prevalence as well as comprehensive prevention measures should be increased [1]. Although prevalent in the endemic areas described, the disease occurs as isolated imported cases in

a number of other countries. Its sidelining from major medical studies is probably due to the fact that it is prevalent mainly in poor, underdeveloped areas of the world and, although it results in severe lower limb deformities, it is rarely fatal.

The parasite *Tunga penetrans* was originally discovered in the Americas. The first empirical descriptions of the biology of the parasite and the changes it causes on the skin of affected patients were described as early as the 16th century by the German Hans Staden [2]. He observed Indian tribes in Brazil and skin changes in them caused by the so-called jigger flea, sand flea or chigo flea. These are several different names that would later be combined into the Latin name *Tunga penetrans*, a pathogenic skin parasite. The first scientific description was in the early 17th century by a Brazilian author who had long observed manifestations of the disease in indigenous tribes [3]. In the 19th century in South America, Africa and the Caribbean, inhabitants of entire settlements began to abandon their habitat and migrate. It appears that the cause was the infection with the jigger, which caused in some of them severe deformities of the lower limbs with ulcers, suppuration, even gangrenous areas [4, 5]. Similar deformities in the lower limbs occurred in some soldiers in the colonial wars [6]. Contemporary authors describe a less dramatic picture, but the continued prevalence of *Tunga penetrans* in poor communities still threatens a number of individuals with the development of severe debilitation. Considering also the increased migration in the world, there is virtually no community that is not threatened by the occurrence of single cases, even in countries with high

economic development and serious disinsection programs [7-9]. *Tunga penetrans* is extremely small in size, about 1 mm. Both female and male fleas feed on blood, but only females permanently penetrate the skin of their hosts, in some cases humans (Figure 2). During penetrance, complex biological changes occur in the flea - its abdomen hypertrophies, abdominal segments grow to about 1 cm, and multiple eggs develop within it. When the eggs are shed, involution of the parasite begins and it dies within a few weeks by detaching from the epidermis, but having accomplished its main goal to lay its eggs in the host [4, 10, 11].

Materials And Methods

In a systematic review and meta-analysis, the group of Obebe et al. established the prevalence and associated risk factors of tungiasis in sub-Saharan Africa (SSA), which is one of the endemic regions in the world [12]. Information on population-based studies was searched in AJOL, Google Scholar, Web of Science and PubMed. SSA data for the period January 1980 and July 2020 were evaluated. Of 27 studies with 16303 individuals, from seven countries in SSA, analyzed in full text, the overall prevalence of disease was 33.4% (95% CI: 27.6-39.8). In the literature sources we adopted, we searched for data on the spread of tungiasis in the mentioned endemic countries, the risk factors for infection related to the socio-economic and hygienic problems of the local population; attention was paid to the possibilities for insecticide prophylaxis and the danger of the spread of tungiasis in non-endemic areas.



Figure 1: Deformed limb from tungiasis (fotos from Michael Wunderli).




Figure 2: *Tunga penetrans*, female (fotos from Michael Wunderli)

Results And Discussion

Ethiopia had the highest prevalence at 46.5% and Uganda the lowest at 20.1%. Risks were associated with sex, age of participants, presence of ground floor (of soil), irregular use of shoes, contact with animals, residence in rural areas. The risk associated with sex varied in most studies and according to most authors the incidence was related to exposure and environmental factors rather than actually to sex [12, 13]. The age-specific prevalence is classical, being highest in children (5-14 years) and adults ≥ 60 years. Children most often work on farms or play on them barefoot, and adults are more likely to overlook or under-see a flea bite because of poor eyesight and lack the ability to remove it from their skin in a timely manner as people of active age [5, 12- 14]. The authors report that the presence of tungiasis can lead to shame and stigma, especially in children, which can limit their attendance at school and socialization [12, 14]. Respectively, prevention measures should also strongly depend on age groups.

The parasite lives 3-5 cm below the sand level as it prefers an oxygen-poor habitat. That is why living in huts with earth floor gives an easy opportunity for infestation. All studies in the field have shown that modification of flooring, usually with cement, severely limits disease incidence [4,12-15]. Some studies have shown that no viable forms of *Tunga penetrans* have been detected in outdoor soil samples, but instead, soil samples under beds and hammocks, as well as in indoor dog kennels, are rich in the parasite (65.0% v. 35.0%; $P=0.02$) [15]. Authors working on the problem describe any part of the body as a possibility of infestation with *Tunga penetrans*, but all are adamant that the feet are most commonly affected due to the fact that the flea hops poorly. Different localisations are possible more commonly in young children playing on the ground [12, 16, 17]. Heukelbach et al. described 1184 residents of a poor neighborhood in Brazil. They found that 33.6% (95% CI : 30.9-36.4) had tungiasis, with only 6% having lesions at sites other than the feet. The second

most frequent location was the upper extremities, especially the elbows, hips and gluteal region [17].

In these areas, walking barefoot is more the norm, and on rare occasions flip-flops are worn. In both cases this has been described as a serious risk factor. Unfortunately wearing shoes in such areas, especially in the absence of such habits, is seen as a physical inconvenience. Children in poor communities consider them a valuable resource and if they own them use them for special occasions - going to school or church for example. Similar evidence is presented for children specifically in Madagascar [10, 12, 14, 18]. Travelling foreigners in such areas mainly wear flip-flops feeling more comfortable in the local rural areas and this is what contributes to their contracting and later transmitting the disease to new communities. In visits to Brazil, Madagascar, Uganda, and Ethiopia between 2007 and 2018, GeoSentinel, as a major global network monitoring travel-related diseases, recorded 87 cases of tungiasis in tourists [19]. In an earlier study covering 1997-2013, 35 tourists from Brazil alone were diagnosed with tungiasis [20].

All authors working on the subject note that the presence of dogs, goats, pigs, cats, in or around the house is an important factor for the development of tungiasis. These animals are usually a serious reservoir for *Tunga penetrans* [6, 8, 21, 22].

An indirect relationship between the incidence of tonsillitis and the type of water supply has also been demonstrated. A link has been made to both family hygiene and the fact that walking barefoot for fresh water for long periods and daily increases the risk of disease [12, 21, 23].

Studies in humans in endemic areas indicate that parasite load is high. Those infected in a study in rural Brazil had an average of 8.9 lesions. The highest numbers were found in males, especially females, children <15 years and very elderly people [10, 23]. In

another study on the prevalence of tungiasis in the Americas for January 2007 to June 2001, PubMed and AJOL databases were used. A total of 83 articles were analyzed, with cases reported in 10 countries, and 71% of cases were in Brazil [24]. In the general population, the prevalence varied between 1.0% and 82.6% depending on several criteria. The age distribution is reported in most publications and it clearly shows that children are the most affected, with a high intensity and severity of the disease [25]. Authors report an S-shaped age curve in statistical studies, with a peak in children 5-12 years and a second peak in adults over 60 years [25, 26].

It is curious how rapid the spread of disease can be among people in resource-poor and poorly hygienic environments. In a study in northeastern Brazil, all individuals who re-entered the community after being out of the endemic area for a long time became infected within about 3 weeks [27]. Studies based in 3 South American countries have shown the importance of seasons. Immediately after the passage of active local rains and the onset of the dry season, the incidence of tungiasis increases about threefold [25- 29]. Remarkably, sick children ≤ 1 year are also found in US communities. Typically, these children are carried on the mother's hips or back and have no contact with soil, and the fact that they are found ill at such a young age speaks to the seriousness of the problem [30].

Conclusion

Over the past 20 years, studies in endemic areas on tungiasis have clarified to some extent the scientific community's knowledge of the epidemiology and risk factors in the Americas. A particular challenge is the fact that *Tunga penetrans* parasitizes a large number of hosts. This makes the parasite highly prevalent in endemic areas. So-called integrated interventions are proposed in this case. These are based on the "One Health" approach and require large-scale cross-sectoral support to reduce DNDi, as those living in these endemic areas often also suffer from a range of other DNDi. This is why urgent measures are needed, such as identifying the places where tungiasis is most intensely prevalent, assessing the severity of symptomatology in relevant areas. It is recommended the development of entomological methods to better understand the transmission of the parasite in different settings and to assess the socioeconomic impact of tungiasis on the community.

References

- World Health Organization. (2020). Ending the neglect to attain the Sustainable Development Goals: A road map for neglected tropical diseases 2021–2030. Geneva: World Health Organization. <https://www.who.int/publications/item/9789240010352>
- Staden von Homberg zu Hessen, H. (1556). *Wahrhaftige Historia und Beschreibung einer Landschaft der wilden, nacketen, grimmigen Menschenfresser Leuten, in der Neuen Welt America gelegen*. Frankfurt am Main: Weigand Hahn. <https://www.academia.edu/RegisterToDownload/RelatedWorks>
- Guerra, F. (1968). Aleixo de Abreu [1568–1630], author of the earliest book on tropical medicine describing amoebiasis, malaria, typhoid fever, scurvy, yellow fever, dracontiasis, trichuriasis and tungiasis in 1623. *Journal of Tropical Medicine and Hygiene*, 71, 55–69. <https://pubmed.ncbi.nlm.nih.gov/4869639/>
- Heukelbach, J., De Olivera, F., & Feldmeier, H. (2003). Ectoparasitoses e saúde pública: desafios para controle. *Cadernos de Saúde Pública*, 19, 1535–1540. <https://www.scielo.br/j/csp/a/LLKQBSqrXMgDnDQ5wHm4FBd/>
- Heukelbach, J., De Olivera, F. A., Hesse, G., & Feldmeier, H. (2001). Tungiasis: A neglected health problem of poor communities. *Tropical Medicine & International Health*, 6, 267–272. <https://pubmed.ncbi.nlm.nih.gov/11348517/>
- Gordon, R. M. (1941). The jigger flea. *The Lancet*, 2, 47–49. [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(00\)71405-X/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(00)71405-X/fulltext)
- Sachse, M., Guldbakke, K., & Khachemoune, A. (2007). Tunga penetrans: A stowaway from around the world. *Journal of the European Academy of Dermatology and Venereology*, 21(1), 11–16. <https://pubmed.ncbi.nlm.nih.gov/17207161/>
- Mukai, Y. (2020). Tunga penetrans in a Sub-Saharan African desert traveler. *Internal Medicine*, 59(19), 2441. <https://doi.org/10.2169/internalmedicine.4652-20>
- Palicelli, A., Boldorini, R., Campisi, P., Disanto, M., Gatti, L., & Portigliotti, L. (2016). Tungiasis in Italy: An imported case of *Tunga penetrans* and review of the literature. *Pathology Research and Practice*, 212(5), 475–483. <https://pubmed.ncbi.nlm.nih.gov/26876782/>
- Eisele, M., Heukelbach, J., Marck, E., Mehlhorn, H., Meckes, O., & Franck, S. (2003). Investigations on the biology, epidemiology, pathology and control of *Tunga penetrans* in Brazil. I. Natural history of tungiasis in man. *Parasitology Research*, 90, 87–99. <https://pubmed.ncbi.nlm.nih.gov/15368123/>
- Heukelbach, J. (2005). Tungiasis. *Revista do Instituto de Medicina Tropical de São Paulo*, 47(6), 307–313. <https://pubmed.ncbi.nlm.nih.gov/16553319/>
- Obebe, O., & Aluko, O. (2020). Epidemiology of tungiasis in Sub-Saharan Africa: A systematic review and meta-analysis. *Pathogens and Global Health*, 114(7), 360–369. <https://pubmed.ncbi.nlm.nih.gov/32909926/>
- Joseph, J. K., Bazile, J., Mutter, J., Shin, S. S., Ruddle, A., & Ivers, L. (2006). Tungiasis in rural Haiti: A community-based response. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 100, 970–974. <https://pubmed.ncbi.nlm.nih.gov/16516941/>
- Muehlen, M., Feldmeier, H., Wilcke, T., Winter, B., & Heukelbach, J. (2006). Identifying risk factors for tungiasis and heavy infestation in a resource-poor community in northeast Brazil. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 100, 371–380. <https://pubmed.ncbi.nlm.nih.gov/16297946/>
- Nagy, N., Abari, E., D'Haese, J., Calheiros, C., Heukelbach, J., & Mencke, N. (2007). Investigations on the life cycle and morphology of *Tunga penetrans* in Brazil. *Parasitology Research*, 101, 233–242. <https://pubmed.ncbi.nlm.nih.gov/17823833/>
- Heukelbach, J., Franck, S., & Feldmeier, H. (2004). Therapy of tungiasis: A double-blinded randomized controlled trial with oral ivermectin. *Memórias do Instituto Oswaldo Cruz*, 99, 873–876. <https://pubmed.ncbi.nlm.nih.gov/15761605/>
- Heukelbach, J., Wilcke, T., Eisele, M., & Feldmeier, H. (2002). Ectopic localization of tungiasis. *American Journal of Tropical Medicine and Hygiene*, 67(2), 214–216. <https://pubmed.ncbi.nlm.nih.gov/4869639/>

- pubmed.ncbi.nlm.nih.gov/12389950/
18. Thielecke, M., Raharimanga, V., Rogier, C., Stauss-Grabo, M., Richard, V., & Feldmeier, H. (2013). Prevention of tungiasis and tungiasis-associated morbidity using the plant-based repellent Zanzarin: A randomized, controlled field study in rural Madagascar. *PLoS Neglected Tropical Diseases*, 7(9), e2426. <https://pubmed.ncbi.nlm.nih.gov/24069481/>
 19. Leder, K., Torresi, J., Libman, M., Cramer, J., Castelli, F., & Schlagenhauf, P. (2013). GeoSentinel surveillance of illness in returned travelers 2007–2011. *Annals of Internal Medicine*, 158(6), 456–468. <https://pubmed.ncbi.nlm.nih.gov/23552375/>
 20. Wilson, M., Chen, L., Han, P., Keystone, J., Cramer, J., & Segurado, A. (2014). Illness in travelers returned from Brazil: The GeoSentinel experience and implications for the 2014 FIFA World Cup and the 2016 Summer Olympics. *Clinical Infectious Diseases*, 58(10), 1347–1356. <https://pubmed.ncbi.nlm.nih.gov/24585698/>
 21. Mutebi, F., Krücken, J., Feldmeier, H., Waiswa, C., Mencke, N., & Sentongo, E. (2015). Animal reservoirs of zoonotic tungiasis in endemic rural villages of Uganda. *PLoS Neglected Tropical Diseases*, 9(10), e0004126. <https://pubmed.ncbi.nlm.nih.gov/26473360/>
 22. Mutebi, F., Krücken, J., Feldmeier, H., Waiswa, C., Mencke, N., & von Samson-Himmelstjerna, G. (2016). Tungiasis-associated morbidity in pigs and dogs in endemic villages of Uganda. *Parasites & Vectors*, 9(44). <https://pubmed.ncbi.nlm.nih.gov/26817587/>
 23. Muehlen, M., Heukelbach, J., Wilcke, T., Winter, B., Mehlhorn, H., & Feldmeier, H. (2003). Investigations on the biology, epidemiology, pathology and control of Tunga penetrans in Brazil. II. Prevalence, parasite load and topographic distribution of lesions in the population of a traditional fishing village. *Parasitology Research*, 90(6), 449–455. <https://pubmed.ncbi.nlm.nih.gov/12768415/>
 24. Saboyá-Díaz, M. I., Nicholls, R., Castellanos, L., & Feldmeier, H. (2022). Current status of the knowledge on the epidemiology of tungiasis in the Americas. *Revista Panamericana de Salud Pública*, 46, e124. <https://pubmed.ncbi.nlm.nih.gov/36060204/>
 25. Feldmeier, H., Kehr, J. D., Poggensee, G., & Heukelbach, J. (2006). High exposure to Tunga penetrans (Linnaeus, 1758) correlates with intensity of infestation. *Memórias do Instituto Oswaldo Cruz*, 101(1), 65–69. <https://pubmed.ncbi.nlm.nih.gov/16612510/>
 26. Vergara, M., Barthel, M., Labarca, M., Neira, O., & Espinoza, R. (2009). Tungiasis afecta a un turista chileno. *Revista Chilena de Infectología*, 26(3), 265–269. <https://pubmed.ncbi.nlm.nih.gov/19621162/>
 27. Heukelbach, J., Franck, S., & Feldmeier, H. (2004). High attack rate of Tunga penetrans (Linnaeus, 1758) infestation in an impoverished Brazilian community. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 98(7), 431–434. <https://doi.org/10.1016/j.trstmh.2003.12.004>
 28. Heukelbach, J., Wilcke, T., Harms, G., & Feldmeier, H. (2005). Seasonal variation of tungiasis in an endemic community. *American Journal of Tropical Medicine and Hygiene*, 72(2), 145–149. <https://pubmed.ncbi.nlm.nih.gov/15741550/>
 29. Pan American Health Organization. (2021). 59th Directing Council: 73rd Session of the Regional Committee of WHO for the Americas. Washington, D.C.: PAHO. <https://www.paho.org/en/documents/cd599-one-health-comprehensive-approach-addressing-health-threats-human-animal>
 30. Muehlen, M., Feldmeier, H., Wilcke, T., Winter, B., & Heukelbach, J. (2006). Identifying risk factors for tungiasis and heavy infestation in a resource-poor community in northeast Brazil. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 100(4), 371–380. <https://pubmed.ncbi.nlm.nih.gov/16297946>