

Kakutani Conjecture

Yang YanHong

Yulong County, Lijiang City, Yunnan Province

*Corresponding author: Yang YanHong, Yulong County, Lijiang City, Yunnan Province,

Submitted: 29 March 2025 Accepted: 03 April 2025 Published: 08 April 2025

Citation: Yang, Y. (2025). Kakutani Conjecture. Wor Jour of Appl Math and Sta, 1(2), 01-02.

Abstract

The Kakutani Conjecture is about performing iterative operations on a number: dividing it by 2 if it is an even number, and multiplying it by 3 and adding 1 if it is an odd number. Will it eventually reach 1? Through the equal quantity transformation $10 = 0$, taking the logarithm \lg , then $1 =$ does not exist. In this proof, a number is written in binary form and then folded in half. By analyzing the structure of the number in the form of 0 and 1 combined with the Zhouyi Bagua (Eight Diagrams), it is concluded that the Kakutani rule is a Mobius strip.

Keywords: Kakutani Conjecture, Kakutani Rule, Life Formula, Equal Quantity Transformation, Zhouyi, Bagua, Qian, Mobius Strip.

Starting from the operation rules of the Kakutani Conjecture, assume that there are numbers $S + 1$ and $S - 1$ that follow the same Kakutani operation rule (S is an integer).

$$\begin{aligned} \text{Then } A &= 3(S + 1) + 1, T = 3(S - 1) + 1 \\ A + T &= 6S + 2 \end{aligned}$$

Starting from the Kakutani operation rule, since $A + T$ is an even number, it should be divided by 2.

Denote the Kakutani operation rule as $L(S) = (A + T)/2 = 3S + 1$

When solving for -5, -7, -17, for $3X + 1$ (the Kakutani operation rule), when the calculation is repeatedly executed, it will enter a cycle. Starting from the operation rules for negative numbers, modify the Kakutani operation rule: for negative odd numbers, repeatedly execute $3X - 1$, and for even numbers, divide by 2.

$$\begin{aligned} G &= 3(S'' + 1) - 1 \\ C &= 3(S'' - 1) - 1 \\ G + C &= 6S'' - 2 \end{aligned}$$

Denote the rule as $F(S'') = (G + C)/2 = 3S'' - 1$.

Then $A + G + T + C = 2*[L(S) + F(S'')]$

Express an integer Y as $Y = \log(N * 1/N * X)$

Then $-Y = -\log(N * 1/N * X)$

$$A + T = 6Y + 2$$

$$G + C = 6(-Y) - 2$$

$$\begin{aligned} L(S) + F(S'') &= 3\log N + 3\log(X/N) + 1 + 3\log N + 3\log[1/(NX)] - 1 \\ &= 6\log N + 3\log(1/N * 1/N) \\ &= 6\log N - 6\log N = 000000 \\ \text{That is, } A + T + G + C &= L(S) + F(S'') \end{aligned}$$

That is, Theorem 1: The property remains unchanged after folding in half.

Write the number S in binary form and then fold it in half. There are four forms corresponding to the starting numbers 0, 1, 10, and 11. Write the Bagua (Eight Diagrams) from the up-down structure form to the left-right structure form $()$, and the corresponding relationship between the 64 hexagrams and the AGCT genetic code can be obtained.

$$\begin{aligned} \text{From } AAA &= AOA \\ \text{Then } x^{\{3\}} &= 2x^{\{2\}} \quad (1) \\ 3x &= 2x + 2 \quad (2) \end{aligned}$$

From the Life Formula (Equal Quantity Transformation)
 $A = T, G = C$
 Then $A + G = T + C$
 That is $x^{\{3\}} + 3x = 2x^{\{2\}} + 2x + 2$
 Then $3x + 1 = 2x^{\{2\}} + 2x + 3 - x^{\{3\}}$
 $X = 2$ makes the equation hold.
 Substitute $x = 3$ into it
 Then $10 = 0$

Check 10, then $10 \rightarrow 5 \rightarrow 16 \rightarrow 8 \rightarrow 4 \rightarrow 2 \rightarrow 1$

That is, after verification, the Kakutani Conjecture also leads to 1 for 10 in the end.

If $x = 3$, we can calculate

$$10 = 0$$

Checkout 10, then $10 \rightarrow 5 \rightarrow 16 \rightarrow 8 \rightarrow 4 \rightarrow 2 \rightarrow 1$

Then: 10 will fall to 1 also.

$$-10 = 0$$

$$-10 \div 2 = -5$$

$$-5 \times 3 + 1 = -14$$

$$-14 \div 2 = -7$$

$$-7 \times 3 + 1 = -20$$

$$-20 \div 2 = -10$$

$$-10 \div 2 = -5$$

For an infinite number ∞ , perform the reciprocal operation

$S = \frac{1}{\infty}$, denoted as $S = 000000$

Since the Bagua Qian in the up-down structure and the Bagua Qian in the left-right structure are equivalent and identical

We can know 000000

Then AOA = AAA

That is $2A^2 = A^3$

$$2A + 2 = 3A$$

Then $A^3 + 3A = 2A^2 + 2A + 2$

$$3A + 1 = 2A^2 + 2A + 3 - A^3$$

When $A = 2$, equation holds.

When $A = 3$, then

$$3A + 1 = 2A^2 + 2A + 3 - A^3$$

$$10 = 0$$

Denoted as $[= O]$

That is, when using a computer to perform the operation of the Kakutani Conjecture, when it comes to an infinite number, there will be a memory overflow, that is, it cannot be verified by a computer.

For example, 20 written in binary is 10100. After folding it in half, there are two cases for the starting number: 1 or 0.

When 20 is written in binary, it can be mathematically expressed as $S = E(s)$

When the starting number is 1, $S = E(s)$ and $s \equiv 2 \pmod{3}$

When the starting number is 0, $s/2$ and $s \equiv 0 \pmod{2}$

Substitute $s = 0$, $s - 1$, $s + 1$ into the Kakutani rule,

We can know that $A = 3(0 + 1) + 1$, $T = 3(0 - 1) + 1$

$$A + T = 2$$

Then there exists the Kakutani rule $L(0) = 1$

That is, write 0 on one side of a paper tape and 1 on the other side, twist and connect them, and we can know that the operation rule of the Kakutani Conjecture is a Mobius strip.

After the Collatz Conjecture converts a number into binary and folds it in half, the starting numbers have four possibilities: 1, 0, 10, 11 (in binary). Assume that after the iterative operation of the Collatz Conjecture, it finally falls back to the starting number. Then from $L(0) = 1$, that is, 0 and 1 are equivalent. Since 10 in binary is equal to 2, we can know that $A = 2$ and equation holds. Since 11 in binary is equal to 3 which is G,

The binary representation of -5 should be expressed in two's complement, which should be 11111011

(The original code is 10000101, the one's complement is 11111010, and the two's complement is 11111011)

The binary folding of -5 represented by symbols is GCGG

Then we can know that $X = -5$ has entered the cycle of $-10 \rightarrow -5 \rightarrow -7 \rightarrow -20 \rightarrow -10$

References

1. "Zhouyi"
2. "Asimov's New Guide to Science", Life Formula